

DNA Barcoding of Marine Metazoa

Ann Bucklin,¹ Dirk Steinke,²
and Leocadio Blanco-Bercial¹

¹Department of Marine Sciences, University of Connecticut, Groton, Connecticut 06340;
email: ann.bucklin@uconn.edu, leocadio@uconn.edu

²Biodiversity Institute of Ontario, University of Guelph, Ontario, N1G 2W1 Canada;
email: dsteinke@uoguelph.ca

Annu. Rev. Mar. Sci. 2011. 3:471–508

First published online as a Review in Advance on
August 18, 2010

The *Annual Review of Marine Science* is online at
marine.annualreviews.org

This article's doi:
10.1146/annurev-marine-120308-080950

Copyright © 2011 by Annual Reviews.
All rights reserved

1941-1405/11/0115-0471\$20.00

Keywords

DNA barcode, marine biodiversity, taxonomy, species identification,
mitochondrial DNA

Abstract

More than 230,000 known species representing 31 metazoan phyla populate the world's oceans. Perhaps another 1,000,000 or more species remain to be discovered. There is reason for concern that species extinctions may outpace discovery, especially in diverse and endangered marine habitats such as coral reefs. DNA barcodes (i.e., short DNA sequences for species recognition and discrimination) are useful tools to accelerate species-level analysis of marine biodiversity and to facilitate conservation efforts. This review focuses on the usual barcode region for metazoans: a ~648 base-pair region of the mitochondrial cytochrome *c* oxidase subunit I (COI) gene. Barcodes have also been used for population genetic and phylogeographic analysis, identification of prey in gut contents, detection of invasive species, forensics, and seafood safety. More controversially, barcodes have been used to delimit species boundaries, reveal cryptic species, and discover new species. Emerging frontiers are the use of barcodes for rapid and increasingly automated biodiversity assessment by high-throughput sequencing, including environmental barcoding and the use of barcodes to detect species for which formal identification or scientific naming may never be possible.

INTRODUCTION

What Is—and Is Not—DNA Barcoding?

DNA barcoding is the derivation of short DNA sequence(s) that enables species identification, recognition, and discovery in a particular domain of life (e.g., Hebert et al. 2003a). The primary purpose of DNA barcoding is to identify an unknown specimen in terms of known classification (Miller 2007, Stoeckle & Hebert 2008), and thus DNA barcoding will complement—not supplant or invalidate—existing taxonomic practices.

It is important to discriminate the goals of DNA barcoding from DNA taxonomy. The latter involves the circumscription and delineation of species based on evolutionary species concepts and uses DNA sequences as the primary and diagnostic characteristics to identify and classify species (Vogler & Monaghan 2006). In contrast, DNA barcoding provides a means of identifying known species by sequence similarity and does not equate species identity, formally or informally, with a particular DNA sequence.

Ideally, barcoding and species discovery can be seen as complementary, with DNA sequences providing diagnostic characters that complement, accelerate, and revive traditional morphological taxonomy but do not replace it (DeSalle et al. 2005). Barcoding studies can help detect overlooked species with subtle or complex morphological traits and stimulate collection of additional genetic, morphological, ecological, geographical, and behavioral data (Bucklin & Frost 2009, Hebert et al. 2004a, Smith et al. 2008b, Steinke et al. 2009b).

The most frequently used gene for metazoans is the mitochondrial cytochrome *c* oxidase subunit I (COI) (Hebert et al. 2003a). The COI barcode region is located at the 5' end of the gene—usually 648 base pairs long—flanked by conserved regions that are the usual basis for design of conserved PCR (polymerase chain reaction) primers, such as the widely used Folmer primers (Folmer et al. 1994). This barcode region has been shown to exhibit a marked divergence between genetic distance within metazoan species (typically <3%) versus that between species (typically 10–25%; Hebert et al. 2003b).

Although the principal of DNA barcoding can be extended to various gene regions, the scope of this review is restricted almost entirely to the COI barcode region usually determined for metazoans. Exceptions are for taxa, described here as “problem children,” for which COI is not a suitable barcode (e.g., corals, sea anemones, and sponges). This review examines the utility (i.e., accuracy, precision, and reliability) of COI for the primary purposes of a DNA barcode in the recognition, discrimination, and discovery of species of metazoans.

History of DNA Barcoding

The basis for DNA barcoding is the decades-long use of DNA sequences in taxonomy and systematics for identifying and classifying organisms. The recent explosion of barcoding interest may be traced to Paul D. N. Hebert from the University of Guelph, Ontario, Canada, who proposed the compilation of a public library of DNA barcodes that would be linked to named specimens (Hebert et al. 2003a). In 2004, the Consortium for the Barcode of Life (CBOL; <http://barcoding.si.edu>) was launched with a mission of developing DNA barcoding as a global standard for the identification of biological species, promoting the rapid compilation of high-quality DNA barcode records in a public library of DNA sequences, and developing new instruments and processes to make barcoding cheaper, faster, and more portable (Savolainen et al. 2005). CBOL now comprises more than 170 member organizations in 50 countries. In partnership with the Census of Marine Life (CoML; <http://www.coml.org>), CBOL recently initiated an international campaign for barcoding marine

biodiversity (MarBOL; <http://www.marinebarcoding.org>). MarBOL is working to ensure that barcodes are available for members of all key marine groups, with good coverage for groups of highest scientific or societal importance.

The extraordinarily rapid advances of DNA barcoding have not been without controversy. The sole use of COI barcodes as characters for species identification was met with initial skepticism (e.g., Will & Rubinoff 2004). Some applications of barcodes have engendered controversy and complaint within the taxonomic community, including using barcodes to delimit species boundaries and reveal cryptic species within known taxa (Wiens 2007), as well as barcode analysis for discovery of new or undescribed species (DeSalle 2006, Rubinoff et al. 2006). An emerging frontier is the construction of barcode libraries for analysis of species biodiversity without the expectation of formal identification and naming of the cataloged species (e.g., Blaxter et al. 2005).

Why DNA Barcoding?

There are an estimated 230,000 described species of marine metazoans and perhaps another 1,000,000 or more species awaiting discovery and description. Our knowledge of life in the oceans is still fragmentary and—for organisms that are tiny, rare, fragile, and/or found in obscure habitats—rudimentary. The diversity of life in the oceans can be analyzed and understood at many levels of organization, including higher taxonomic levels, functional groups, species, populations, and individuals. The careful process of collecting, classifying, and identifying known species and describing novel species will continue for decades to come. In the meantime, new methodologies are needed to expedite and facilitate the work of documenting current biodiversity levels and distribution patterns in the ocean.

Marine species may have extensive biogeographic ranges, spanning multiple ocean basins and reaching from pole to pole. Oceanographers and marine biologists are still working to map the oceans and describe the diversity, distribution, and abundance of metazoan species. Marine biodiversity can be examined using morphological characters and classical taxonomy, molecular systematic approaches (e.g., DNA taxonomy), and various other remote observation technologies, such as optical and acoustical imaging. Yet the recognition and discrimination of species is one of the foundations of biodiversity research, and it is exactly this step that is the most time-consuming and difficult—frequently forming a bottleneck in the analytical chain of sample and data collection and analysis.

DNA sequence analysis offers many opportunities for accurate, reliable, rapid, and eventually remote identification of specimens of all metazoan groups. DNA barcodes may be particularly useful for species identification of organisms that are rare, fragile, and/or small, especially when morphological identification is difficult and mistakes are likely due to simple or evolutionarily conserved body plans. Also, many taxa have circumglobal or disjunct geographic distributions; barcodes can reveal taxonomically significant geographic variation and cryptic species (see Ten Reasons for Barcoding sidebar, below).

DNA barcode libraries will lead to rapid molecularly based analysis of samples for known species. DNA barcodes are tools for understanding species diversity for global ocean/earth observing initiatives. Neither morphology nor genetics should be oversimplified; both can assess variation within and between populations and species, evolutionary relationships among species and higher taxa, and processes associated with speciation. Much may be lost in translation, but the goals of morphologists and geneticists studying marine organisms are the same: to gain genuine knowledge of species diversity of life in the ocean.

TEN REASONS FOR BARCODING^a

1. Works with fragments. Barcoding can identify a species from bits and pieces. Once established, barcoding will quickly identify undesirable material in processed foodstuffs and detect commercial products derived from regulated species (e.g., protected whale species served as sushi).
2. Works for all stages of life. Barcoding can identify a species in its many forms (e.g., eggs and larvae of marine fishes).
3. Unmasks look-alikes. Barcoding can distinguish among species that look alike, uncovering dangerous organisms masquerading as harmless ones and enabling a more accurate view of biodiversity (e.g., sibling species swarms of planktonic copepods).
4. Reduces ambiguity. Written as a sequence of four discrete nucleotides—CATG—along a uniform locality on genomes, a barcode of life provides a digital identifying feature, supplementing the more analog gradations of words, shapes, and colors. A library of digital barcodes will provide an unambiguous reference that will facilitate identifying species invading and retreating across the ocean and through centuries.
5. Makes expertise go further. The bewildering diversity of approximately 230,000 known species of 31 phyla confines even an expert to morphological identification of only a small portion of marine metazoa. Foreseeing perhaps 1,000,000 more species to go, scientists can equip themselves with barcoding to speed identification of known organisms and facilitate rapid recognition of new species.
6. Democratizes access. A standardized library of barcodes will empower many more people to call by name the marine species they observe. It will make possible identification of species abundant and rare, native and invasive, and engender appreciation of marine biodiversity locally and globally.
7. Opens the way for an electronic handheld field guide. Barcoding links biological identification to advancing frontiers in DNA sequencing, miniaturization in electronics, and computerized information storage. Integrating those links will lead to portable desktop devices and ultimately to handheld barcoders.
8. Sprouts new leaves on the tree of life. Barcoding the similarities and differences among marine metazoan species already named will provide a wealth of genetic detail, helping to draw the tree of life on Earth.
9. Demonstrates the value of collections. Compiling the library of barcodes begins with the millions of specimens in museums, aquaria, and other biological repositories. Barcoding shines a spotlight on these institutions and their collections and will strengthen their ongoing efforts to preserve marine biodiversity.
10. Speeds writing the encyclopedia of life. Compiling a library of barcodes linked to vouchered specimens and their binomial names will enhance public access to biological knowledge, helping to create an online encyclopedia of marine animals, with a Web page for every species.

^aModified from the Consortium for the Barcode of Life (<http://barcoding.si.edu>).

Why Mitochondrial Cytochrome C Oxidase Subunit I?

Today's gold standard metazoan barcode is a 658–base pair region at the 5' end of the mitochondrial COI gene (Hebert et al. 2003a). In the search for a universal metazoan barcode gene, the mitochondrial genome of metazoans was considered to have numerous advantages over the nuclear genome: lack of introns, limited exposure to recombination, high copy numbers in every cell, haploid character, and a generally strict maternal mode of inheritance. Many previous studies have used mitochondrial 12S and 16S rRNA genes for systematic analysis, but they share a disadvantage in the prevalence of insertions and deletions (indels) that greatly complicates sequence alignment and comparative analysis. The 13 protein coding genes in the metazoan mitochondrial genome generally lack indels, as these would result in codon reading frame shift mutations. COI was chosen over other mitochondrial protein coding genes because (a) conserved “universal” primers for




Figure 1

Diagrammatic hypothetical frequency distributions of genetic distances within and between species for two cases: (a) the barcode gap (Meyer & Paulay 2005) formed by the discrete distributions of intraspecific variation (shown in red) and interspecific variation (shown in yellow), and (b) overlapping distributions between intra- and interspecific variation and no barcode gap. Figure reproduced from Meyer & Paulay (2005).

a sizable region of the variable 5' end of the gene had proved useful for species from a broad range of metazoan phyla (Folmer et al. 1994), and (b) COI, compared with other mitochondrial genes, appeared to possess a phylogenetic signal useful over a broader range of taxonomic levels. Similar to other protein coding genes, third codon position nucleotides show a high incidence of base substitutions, leading to a rate of molecular evolution nearly three times greater than that of mitochondrial rRNA genes. Importantly, COI evolution is sufficiently rapid to allow the discrimination of very closely related species in most groups, as well as taxonomically significant intraspecific variation associated with geographic structure.

The primary reason for the selection of COI as the gold standard barcode gene is the typical pattern of variation observed for numerous species, with marked divergence and lack of overlap between intraspecific (i.e., between individuals of the same species) and interspecific (i.e., between individuals of different species) genetic distances (Hebert et al. 2003b). Dubbed the “barcode gap” by Meyer & Paulay (2005) (Figure 1), the lack of overlap between intra- and interspecific variation has been deemed of paramount importance for the accuracy and reliability of barcode genes. Consistent findings through many studies of diverse metazoans, including analysis of intraspecific variation of geographically widespread species, have led to the adoption of COI as the gold standard marker for the barcoding community (Stoeckle & Hebert 2008). This choice is clearly reflected in the large public databases such as the National Center for Biotechnology Information (NCBI) GenBank database (<http://www.ncbi.nlm.nih.gov/genbank>) and the Barcode of Life Data Systems (BOLD; <http://www.boldsystems.org>).

Exceptions to the resolving power of COI and other mitochondrial genes include the phyla Porifera (sponges) and Ctenophora (comb jellies) and the cnidarian class Anthozoa (corals and sea

anemones). The evolutionary rate of mitochondrial DNA (mtDNA) of these animals was found to be 10–20 times lower than other metazoans (Shearer et al. 2002, Huang et al. 2008) and usually insufficient to allow reliable discrimination between closely related species.

Standardization has resulted in a flood of COI sequences into public repositories such as BOLD and GenBank. The exceptional diversity and density of the data have allowed examination of molecular evolutionary hypotheses, including evolutionary changes in nucleotide base composition, such as GC shifts in animals (Min & Hickey 2007), as well as plants and fungi (Clare et al. 2008). Although the COI barcode region is not an especially effective sentinel sequence in predicting nucleotide composition (and likely no different from any region of the mitochondrial genome), barcode libraries will likely contain many replicates, with dense sampling across diverse taxa and thus offer many opportunities for analysis not possible for more limited mitochondrial whole-genome databases.

Other Genes

Numerous studies have compared COI and other gene sequences in terms of usual barcode functions: identifying species, discriminating closely related species, revealing cryptic species, and discovering new species. Erpenbeck et al. (2006) suggested the inclusion of another partition of COI (I3-M11, 23 base pairs downstream of the barcode region) to enable discrimination of species of cnidarians. The internal transcribed spacers (ITS) of nuclear rRNA genes have been evaluated as barcodes for many marine groups, especially those for which COI has proved problematical. A review (Vollmer & Palumbi 2004) of more than 500 studies published in 2000–2004 that used ITS as the main marker found many reports of intraindividual variability, suggesting that caution is required when analyzing these sequences. Calderón et al. (2006) compared patterns of COI and ITS variation within and between species of gorgonian corals in the Mediterranean; the gene regions used both identified species but did not resolve population structure. Chu et al. (2009) compared COI, mitochondrial 16S rRNA, and nuclear 18S rRNA sequences as barcodes for birds, fishes, nematodes, and crustaceans; they concluded that 18S barcodes were advantageous for some analytical methods. Whether or not the COI barcode region alone is sufficient for species assignments—a sometimes contentious issue—it is widely viewed as advantageous to use sequences from multiple loci (Neigel et al. 2007).

BIOINFORMATICS AND STATISTICAL ANALYSIS

The bioinformatics of DNA barcoding can be divided into two distinct approaches: (*a*) diagnostic identification of species (i.e., the correct assignment of an unknown sample to a known species), and (*b*) detection and classification of novel sequences for previously unsampled species. Diagnostic identification requires that an individual's DNA sequence can be unambiguously associated with the sequences of a group of organisms, ideally at the taxonomic level of species. Establishing this association is complicated for several reasons, including patterns of intraspecific variation of the reference sequence and possible incongruence between the evolutionary histories of the gene and species.

Most recently published studies have used analysis of genetic distances to infer the species identification based on the barcode designation. One approach is a simple NCBI BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) query (Altschul et al. 1990), which uses a raw similarity score to determine the nearest neighbor to the query sequence. Although BLAST searches are rapid, they have a number of significant drawbacks, including high rates of false-positive results

inherent to the BLAST algorithm and frequent sampling density-dependent accuracy (Koski & Golding 2001).

Other methods such as TaxI (Steinke et al. 2005) and TaxonDNA (Meier et al. 2006) apply linear searches based on distances of pairwise comparisons to infer the most similar sequence and identify the species based on the barcode sequence.

BOLD was initially developed as an informatics workbench for a single, high-volume DNA barcode facility (Ratnasingham & Hebert 2007). BOLD uses an identification (ID) engine that collects nearest neighbors from all reference sequences by aligning the query to a global alignment through a Hidden Markov Model (HMM) profile of the COI protein (Eddy 1998), followed by a linear search of the reference library. BOLD, currently the most commonly used species ID system based on barcodes, has evolved into a community resource and is used by CBOL barcoding campaigns for fishes and birds, as well as MarBOL.

It should be noted that the accuracy of the BOLD algorithm (and all other distance-based methods) depends on the degree of disparity between intra- and interspecific sequence variation (i.e., the barcode gap discussed above). Whereas the existence of such a gap has been successfully demonstrated in a variety of studies (Hajibabaei et al. 2006a, Kerr et al. 2007, Ward et al. 2005), other studies have shown that young species may lack a distinctive barcode gap (e.g., up to 8% of the gastropod group cowries; Meyer & Paulay 2005). Furthermore, the analysis of species affiliation and the inference of species identity require an a priori threshold similarity value; it remains questionable whether universal thresholds exist, even among congeneric species (Ferguson 2002, Hickerson et al. 2006, Whitworth et al. 2007).

Recently, statistically more sophisticated methods of species identification have been developed. Matz & Nielsen (2005) described a likelihood-ratio test (LRT) to query species identity. Nielsen & Matz (2006) and Abdo & Golding (2007) used coalescent-based approaches in the context of Markov Chain Monte Carlo (MCMC) methods to assign unknowns to identified species. All these methods are computationally demanding, and in the case of the LRT, corrections for multiple testing remain uncertain (Ross et al. 2008). Furthermore, the Bayesian method currently implemented (Nielsen & Matz 2006) cannot handle more than two species at a time, thereby limiting its use for DNA barcoding. Although the decision-theory method (Abdo & Golding 2007) uses more information in the sequence data than distance-based methods, the statistical power comes with a huge computational expense, limiting its usefulness. None of these methods is a panacea, yet each offers potential improvement over previously used methods.

Character-based methods have been proposed to identify classification rules based on an existing hierarchical organization and then to classify unknown data (DeSalle et al. 2005, Sarkar et al. 2008). A first application is based on the Characteristic Attributes Organization System (CAOS) algorithm developed for discovery of conserved character states from cladograms or groups of categorical information (Sarkar et al. 2002a,b). Although this method has proved to be fast, accurate, and sufficiently sensitive to distinguish sibling species pairs (Kelly et al. 2007, Lowenstein et al. 2009, Rach et al. 2008), it has not been tested with very large and diverse data sets, such as GenBank and BOLD. CAOS avoids some problems of distance-based methods, such as the reliance on a barcoding gap (Meyer & Paulay 2005, Moritz & Cicero 2004) and is thought by some researchers to be compatible with classical character-based taxonomy (DeSalle 2005, Rubinoff et al. 2006). Another character-based approach adopted a logic mining method based on two optimization models; Bertolazzi et al. (2009) reported results for two data sets where COI barcodes were used to identify individuals to different species.

A recent method by Zhang et al. (2008) is based on existing back-propagation neural networks and uses both differences between sequences and their patterns (e.g., relative positions of variable sites). Neural networks have the disadvantage that they are governed by supervised training and

the parameter(s) responsible for an assignment may remain unknown. Also, input sequences must be assigned to a known species, requiring a complete set of predefined taxa, so the method is not useful for uncertain species identifications or species discovery.

Indicator vectors have been used by Sirovich et al. (2009, 2010) to assign sequences to known species and groups of organisms. Sequences are transformed into vectors; species- and group-distinguishing vectors are determined by contrast with other vectors. The results are presented in visually intuitive scalable displays, called Klee diagrams because of their resemblance to modern art (**Figure 2**).

The second bioinformatic approach of barcoding is the classification of new data based on the distribution of sequence variation within and between species. A specific goal of this approach is to determine whether barcodes can be used to identify previously unrecognized species. The proposed mechanism for the evaluation of unknowns within a partially sampled phylogeny is through the implementation of thresholds, chosen to separate intraspecific variation from interspecific differences. Thus, if two specimens putatively belonging to the same species diverge by ten times the average divergence of that group, then one specimen is likely to be a new species (Hebert et al. 2004b) that warrants further taxonomic examination. This rule may be a useful guide but will detect cryptic species only if they are quite divergent from recognized species (Hickerson et al. 2006, Ward 2009). Currently, classification of barcodes is usually done using a clustering algorithm (e.g., distance-based Neighbor Joining, or NJ) that evaluates genetic distances within and between species (Bucklin et al. 2007, Steinke et al. 2009b, Ward et al. 2005). The limits to this method include undersampling due to poor taxonomy (Meyer & Paulay 2005), ancestral polymorphism, and incomplete lineage sorting. Also, mtDNA sequence divergence does not necessarily track species boundaries (Avice 2000, 2004). For these reasons, some studies have used nuclear markers to corroborate barcode analyses (Bucklin & Frost 2009, Raupach et al. 2009, Smith et al. 2007, Sonnenberg et al. 2007).

DNA barcoding analytical methods, algorithms, and bioinformatic approaches have not yet come of age. Distance-based methods are now widely used to assign an unknown specimen to a known species, detect novel sequences, and determine an unknown specimen to be a distinct species. They are computationally fast and yield consistent results for the many species that exhibit the necessary disparity between intra- and interspecific variation. The barcoding community increasingly acknowledges that distance-based approaches are inconsistent with classical character-based taxonomy (DeSalle et al. 2005) and that valuable information is lost by converting sequences into genetic distances (Zhang et al. 2008). New algorithms and methods are in development that represent the next generation of barcode analysis for both correct assignment of barcodes to known species and classification of novel sequences.

PROGRESS IN BARCODING MARINE METAZOA

The numerous and diverse metazoan phyla found in marine environments provide an excellent opportunity to examine the broad taxonomic utility of COI barcodes for species identification, discrimination, and discovery. We report here a sampling of the barcoding studies that have used COI to examine important taxonomic groups of marine metazoa (**Table 1, Figure 3**).

Taxon-by-Taxon Barcoding

Sponges. The lack of complex morphological characters and high degree of homoplasy have confounded classification within the phylum Porifera (sponges), making DNA barcoding particularly useful in this group. Sponges are highly speciose, ecologically important, and




Figure 2

Vector analysis of 4,176 barcodes of 367 species of elasmobranchs (chimaeras, sharks, and rays) using the methods of Sirovich et al. (2009, 2010). Only species with three or more individuals were used. Results are shown as a Klee diagram, so-named for the resemblance to modern art. (a) Major elasmobranch groups are clearly defined by high correlation coefficients from the vector analysis. (b) Enlargement of a portion of the diagram showing families and genera of rays with highly correlated barcodes and demonstrating the zoom function of vector analysis and Klee diagrams. (Unpublished data and figure from D. Steinke, Univ. of Guelph, Canada.)

commercially valuable to the pharmaceutical and biomaterials industries. The Sponge Barcoding Project (Wörheide & Erpenbeck 2007) aims to examine all species. To date, barcoding has worked reasonably well (Blanquer & Uriz 2007, Erpenbeck et al. 2007, Raleigh et al. 2007). However, the resolving power of COI in sponges appears not to correspond to their higher metazoan counterparts (Cardenas et al. 2009, Heim et al. 2007, Park et al. 2007, Wörheide 2006).

Table 1 Progress in barcoding marine metazoa by phylum^a

Phylum	No. of known species	No. of barcoded species	Barcoded species (%)
Acanthocephala	600	10	1.7
Annelida	12,148	635	5.2
Arthropoda	47,217	3,580	7.6
Brachiopoda	550	35	6.4
Bryozoa	5,700	20	0.4
Chaetognatha	121	23	19.0
Chordata	21,517	7,279	33.8
Cnidaria	9,795	594	6.1
Ctenophora	166	0	0.0
Cycliophora	1	1	100.0
Echinodermata	7,000	771	11.0
Echiura	176	2	1.1
Entoprocta	170	0	0.0
Gastrotricha	400	0	0.0
Gnathostomulida	97	8	8.2
Hemichordata	106	2	1.9
Kinorhyncha	130	0	0.0
Loricifera	18	0	0.0
Mollusca	52,525	4,813	9.2
Nematoda	12,000	180	1.5
Nematomorpha	5	0	0.0
Nemertina	1,230	81	6.6
Orthonectida	24	0	0.0
Phoronida	10	0	0.0
Platyhelminthes	15,000	124	0.8
Porifera	5,500	67	1.2
Priapulida	8	1	12.5
Rhombozoa	82	0	0.0
Rotifera	50	20	40.0
Sipuncula	144	15	10.4
Tardigrada	212	9	4.2
TOTALS	192,702	18,270	9.5

^aNumbers of known species are from Bouchet (2006). Numbers of barcoded species are as reported in the National Center for Biotechnology Information (NCBI) GenBank and the Barcode of Life Data Systems (BOLD) databases. Percentages of barcoded species are determined from numbers as shown.

COI divergence rates are 10–20 times lower for sponges than for Bilateria (Shearer et al. 2002, Wörheide 2006), and the barcode region typically does not distinguish between species. Some studies have used alternative barcode regions, such as the I3-M11 region of COI (Erpenbeck et al. 2006) and ITS (Park et al. 2007, Pöppe et al. 2010).

Cnidarians. Very low levels of COI polymorphism within the cnidarian class Anthozoa (including sea anemones, corals, and sea pens) have meant that the barcode region is of limited usefulness




Figure 3

Numbers of known and barcoded marine metazoan species per phylum. The phyla are ordered by relative numbers of marine species; the nine most speciose phyla (*top*) contain 95% of all marine metazoan species. Blue bars indicate numbers of known species; red bars indicate numbers of barcoded species. The scale is logarithmic. See **Table 1** for actual numbers. Numbers of known species are based on data from Bouchet (2006).

for identifying or discriminating species of this group when they are relatively closely related (Shearer et al. 2002). The underlying evolutionary dynamics may include very slow nucleotide substitution rates for mitochondrial genes, although nuclear genes may vary (Hellberg 2006). Shearer & Coffroth (2008) suggest that the lack of COI divergence among many species indicates the need for alternate markers. There are exceptions to this general conclusion, however. COI discriminated four species of gorgonian corals in the Mediterranean (Calderón et al. 2006). COI discriminated species of the hexacorallian order Zoantharia (Sinniger et al. 2008); genetic distances differed among species, genera, and families (Sinniger et al. 2010), indicating that COI may be useful for zoanthid classification.

COI evolution in the cnidarian classes Hydrozoa and Scyphozoa is more typical of other metazoans (Huang et al. 2008). Multiple cryptic hydrozoan species were discovered based on COI by Folino-Rorem et al. (2009) and Moura et al. (2008). In concert with other genes, COI was used to resolve species and systematic relationships of the hydrozoan family

Campanulariidae (Govindarajan et al. 2006). Within the Scyphozoa, COI has been widely used as a barcode to reveal cryptic species, confirm species identification, and examine evolutionary relationships (Dawson 2005a,b,c; Dawson & Jacobs 2001; Holland et al. 2004). Across the Medusozoa (a cnidarian subphylum comprising of ~3,800 species in the classes Hydrozoa, Cubozoa, and Scyphozoa), COI was shown by Ortman et al. (2010) to be broadly useful as a barcode for species identification and discovery.

Barnacles. Barcoding studies of barnacles in the western Pacific have provided evidence of cryptic species within the acorn barnacle *Chthamalus* (Tsang et al. 2008) and the symbiotic coral barnacle *Wanella* (Tsang et al. 2009) as well as lack of genetic divergence between species of *Tetrachita* (Tsang et al. 2007).

Copepods. Numerous barcoding studies have sought to reexamine morphologically based descriptions of copepod sibling species swarms, which present persistent challenges to expert taxonomists. Among the calanoid copepods, the COI barcode region has been determined and confirmed to be diagnostic for sibling species of *Calanus* (Hill et al. 2001, Unal et al. 2006), *Clausocalanus* (Bucklin & Frost 2009), *Neocalanus* (Machida et al. 2006), and *Pseudocalanus* (Bucklin et al. 2003). Among cyclopoid copepods, species of Oncaeidae—with more than 100 species and many sibling species, a challenge for taxonomists—were examined for genetic distinctiveness of COI and other genes by Böttger-Schnack & Machida (2010). Although copepods are frequently thought of as taxonomically well known, comprehensive analyses based on DNA barcodes are revealing considerable hidden diversity and cryptic speciation across the suborder. Dippenaar et al. (2010) examined the cosmopolitan elasmobranch symbiotic copepod *Nesippus orientalis* (Siphonostomatoidea) off South African coasts and found two divergent clades related to host species; they expect further discoveries of cryptic species within this group. DNA barcoding of another siphonostomatoid copepod, the common sea louse or parasite of fish, revealed species relationships and evidence of cryptic species (Øines & Heuch 2005).

Stomatopods. Barcodes have been used to identify stomatopods (mantis shrimp) from the South China Sea by Tang et al. (2010), who successfully matched unknown larvae to known adult species from the region. In contrast, Barber & Boyce (2006) used barcodes to discriminate larval stomatopods in the Coral Triangle and the Red Sea, but few could be successfully matched to known adults. These results demonstrated the marked underestimation of biodiversity in this well-studied group and highlighted taxonomic limitations of the reference sequence database.

Mysids. Remerie et al. (2006) found evidence of a complex of cryptic species within the mysid *Mesopodopsis slabberi*, with marked divergences between Atlantic and Mediterranean populations.

Amphipods. Analysis of systematic relationships within the amphipod genus *Gammarus*—a highly speciose genus found to be monophyletic using COI and other molecular markers (Hou et al. 2007)—has revealed strong evidence based on the COI barcode region of geographic divergence (Krebs et al. 2010) and cryptic speciation (Costa et al. 2009, Hou et al. 2009). Havermans et al. (2010) used COI to examine the systematics of Antarctic lysianassoid amphipods, finding incongruence between the molecular phylogeny and morphological classification. Radulovici et al. (2009) found cryptic and invasive amphipod species in the Estuary and Gulf of St. Lawrence, in the northwest Atlantic Ocean.

Euphausiids. Frequently considered to be easily recognizable, euphausiids are ecologically important and may occasionally dominate pelagic communities in terms of abundance and biomass. DNA barcodes for euphausiids are useful for species identification, especially of larval and juvenile forms that can be genuinely cryptic. COI reliably discriminated 40 species of euphausiids in a barcoding study by Bucklin et al. (2007). COI proved useful to examine species-specific evolutionary patterns for *Nyctiphanes* (D'Amato et al. 2008).

Decapods. Many decapods are clearly distinguishable as adults but have larval and juvenile forms that are difficult—or impossible—to identify to species. In some instances, the larvae are distinguishable but not easily matched to the correct adult form. New species descriptions of the East Pacific galatheid squat lobster *Munidopsis* included DNA barcodes, which also revealed high gene flow among distant conspecific populations (Jones & Macpherson 2007). Barcodes clearly discriminated species of the fiddler crab *Uca* in the northwestern Indian Ocean (Shih et al. 2009) and the hermit crab *Clibanarius* of the Ryukyu Archipelago of Japan (Hirose et al. 2010). Although barcodes for coral reef decapods could not be matched to known sequences in the GenBank database, they could be used to count numbers of species (Plaisance et al. 2009). A barcoding study of megalops larvae of *Cancer* crabs in the southeastern Pacific Ocean resulted in the description of diagnostic morphological characters for species identification (Pardo et al. 2009). Phylogeographic analysis of decapods using COI has revealed cryptic species within the brachyuran mangrove crab *Perisesarma guttatum* on the eastern coastline of Africa (Silva et al. 2010).

Mollusks. DNA barcoding has been widely used for marine mollusks, including chitons (Kelly et al. 2007), gastropods (Hunt et al. 2010; Jennings et al. 2010a; Krug et al. 2007; Meyer & Paulay 2005; Puillandre et al. 2009a,b,c), bivalves (Feng et al. 2010, Lorion et al. 2009, Mikkelsen et al. 2007, Nuryanto et al. 2007), and cephalopods (Allcock et al. 2010, Undheim et al. 2010). Barcodes discriminated frequently confused species and revealed species relationships of the gastropod *Nerita* throughout the southwestern tropical Pacific Ocean (Spencer et al. 2007). Johnson et al. (2008) found 14 nominal species and numerous cryptic species of deep-sea lepetodrilid limpets, the most abundant and diverse gastropod group at hydrothermal vents. Barcodes have also revealed lack of genetic differentiation among some species of mollusks: sympatric intertidal limpets (Siphonariidae) off coastal southeast Africa lacked barcode differences, suggesting that they are morphotypes of a single species (Teske et al. 2007b). Two clams of the genus *Donax* showed no significant barcode variation and were found to represent one species (Carstensen et al. 2009). Puillandre et al. (2009c) clearly demonstrated the ability of barcodes to identify gastropod larvae, although barcode data are sparse and taxonomic coverage is biased toward shallow water species. Meyer & Paulay (2005) also emphasized problems associated with limited sampling in delineating closely related species in taxonomically understudied groups.

Annelids. The vast majority of marine annelids belong to the class Polychaeta. The group is diverse, with over 12,000 known species and undoubtedly many thousands of species awaiting discovery and description. In a comprehensive barcoding study of polychaetes in the Alaskan and Canadian Arctic, Mincks Hardy et al. (2010) estimated that one-fourth of polychaete species were composed of two or more distinct genetic lineages. A second study found cryptic species within the cosmopolitan fireworm *Eurythoe complanata* (Barroso et al. 2010). Similarly, Rice et al. (2008) found a cryptic species complex within *Polydora cornuta* along the Atlantic and Pacific coasts of North America; barcodes revealed cryptic species within the common ragworm *Hediste diversicolor* in the Baltic Sea (Virgilio et al. 2009). COI discriminated closely related sympatric species of the

lugworm *Arenicola* (Luttikhuizen & Dekker 2010) and the oligochaete *Tubificoides* (Erséus & Kvist 2007) in northeast Atlantic and Scandinavian waters.

Echinoderms. Barcodes discriminated 191 species of echinoderms, including representatives of all five classes (ophiuroids, asteroids, echinoids, holothuroids, and crinoids), confirming the usefulness of sequence variation for species discrimination across the phylum (Ward et al. 2008). Barcoding studies have used COI to examine species limits and reveal cryptic species of asteroids (Naughton & O'Hara 2009), holothuroids (Uthicke et al. 2010), and crinoids (Helgen & Rouse 2006, Wilson et al. 2007). Coordinated efforts to accelerate barcoding of echinoderms have yielded new protocols and primers for COI sequencing (Hoareau & Boissin 2010).

Urochordates. Barcodes have allowed accurate species identification for ascideans (sea squirts) that frequently lack diagnostic morphological characters. Pérez-Portela et al. (2007) found cryptic species within the ascidean *Pycnoclavella* in Atlantic and Mediterranean regions; barcodes were used by Hirose & Hirose (2009) to identify species of *Diplosoma*. Pérez-Portela & Turon (2008b) found that COI resolved relationships among ascidean species and genera but not families (Pérez-Portela et al. 2009).

Flatworms. Vilas et al. (2005) used COI for discovery of cryptic species of parasitic flatworms (phylum Platyhelminthes). COI also revealed cryptic species of a free-living flatworm (Sanna et al. 2009). The community is still in search of optimal marker genes and protocols for a broad barcoding effort (Moszczyńska et al. 2009).

Nematodes. Morphological identification of individual nematodes to described species is often not possible due to their sheer abundance, small size, and lack of taxonomic experts. Nematode diversity in marine sediments remains essentially unknown; surveys suggest that diversity of marine nematodes may exceed 1,000,000 species (Lambshhead 2001). Difficulties in developing COI primers and protocols resulted in the use of the small-subunit (18S) rRNA gene for species identification and diversity estimates (Bhadury & Austen 2006, 2010; Floyd et al. 2005). Recently work has also focused on COI for barcoding (Cross et al. 2007, Elsasser et al. 2009, Ferri et al. 2009). COI variation of the nematode *Thoracostoma trachygaster* revealed cryptic diversity and population genetic structure consistent with dispersal scales of a few 100 km (Derycke et al. 2010). Another study of *Monocelis lineata* showed marked genetic differentiation between the Atlantic and Mediterranean regions and supported the existence a cryptic sibling species complex (Sanna et al. 2009).

Nemerteans. For some taxonomically challenging groups, barcodes alone cannot resolve the systematic questions. Sundberg et al. (2010) barcoded species of the nemertean *Cerebratulus* and were unable to uniquely resolve questions of species identity and relationships based upon the COI barcode region.

Bryozoa. Gomez et al. (2007b) found close correspondence between barcode lineages and reproductive isolation of taxa, despite minimal morphological change and noted the divergence of the cosmopolitan bryozoan *Celleporella* into four major clades (Gomez et al. 2007a).

Pycnogonids. Cryptic lineages within the pycnogonid *Colossendeis megalonyx* were found through COI analysis of Antarctic and sub-Antarctic collections (Krabbe et al. 2010).

Chaetognaths. Subtle morphological differences divide chaetognath species, many of which have extensive biogeographical ranges. Accurate species identification is possible using the COI barcode region, despite high levels of intraspecific variation perhaps resulting from large-scale geographic structuring (Jennings et al. 2010b). Miyamoto et al. (2010) found cryptic species within the deep-sea chaetognath *Caecosagitta macrocephala*.

Fishes. Fishes currently represent the most comprehensively sampled and studied group of marine metazoans. The need for comprehensive and reliable species identification tools, combined with early barcoding success with fishes (Savolainen et al. 2005, Ward et al. 2005), led to the formation of the global initiative Fish Barcode of Life (FISH-BOL; <http://www.fishbol.org>). Launched in 2005, FISH-BOL is gathering DNA barcode records for all 30,000 known species of fishes. To date, barcoding has been completed for 51,906 specimens of 7,437 species, most of which are marine. The barcodes are being used to address issues related to commercial fisheries (e.g., market substitution and quota management) and sampling has been guided by these concerns. FISH-BOL also promises to be a powerful tool for extending the understanding of the natural history and ecological interactions of fish species (Ward et al. 2009).

Recent efforts have developed and optimized analytical protocols for fishes, including DNA extraction and PCR amplification for high-quality barcode sequence data (Ivanova et al. 2006, 2007). Group-specific PCR primers (Paine et al. 2007, Ward et al. 2005) and primer cocktails (Ivanova et al. 2007) are typically used for barcoding fishes. In a few instances (e.g., some elasmobranchs and muraenids), more taxonomically specific primers are necessary (Spies et al. 2006)

DNA barcoding can be used to identify whole fish, as well as fillets, fins, fragments, larvae, and eggs. For all but the whole fish, molecular analysis is particularly useful because the morphological traits used for species identification are generally absent. DNA barcoding was used to identify shark fins confiscated from illegal fishers in northern Australian waters (Holmes et al. 2009), based on a comprehensive barcode library assembled in previous years (Ward et al. 2007, 2008). COI barcoding has been successfully used to identify fish larvae taken from the Great Barrier Reef (Pegg et al. 2006), Caribbean (Victor 2007), and Pacific (Hubert et al. 2010, Paine et al. 2008).

The first fish barcode study (Ward et al. 2005) examined a number of taxa that had only gained provisional recognition from morphological examination. In every case, likely species were separable by their barcodes; some have subsequently been named. Species of the dogfish initially referred to as *Squalus* sp. B, C, D, E, and F have now gained formal scientific names (Last et al. 2007b, Ward et al. 2007). A study of fish species thought to bridge South African and Australian waters found that approximately one-third of those species actually represented two taxa (Zemlak et al. 2009). There are numerous examples among fishes of the power of an integrated approach to taxonomy using barcoding. The first vertebrate to include a COI barcode as part of its species description was the goby *Coryphopterus kuna* (Victor 2007). COI barcoding helped validate a number of new fish species, including a goby (Victor 2008), sting ray, Antarctic ray *Bathyraja* (Smith et al. 2008b), handfish *Brachionichthys australis* (Last et al. 2007a), and five new species of damselfish *Chromis* (Pyle et al. 2008).

Larger-scale barcoding studies of fishes of different regions revealed an average 2% overlooked species based on the distance metric proposed by Hebert et al. (2004b), which targets for further study any barcode that diverges from its putative species by ten times the average divergence within that group. These flagged instances have been subsequently validated by morphology, geography, and ecology and are awaiting formal description (**Table 2**). Extrapolating species discoveries to date among 2% of known species, we can expect up to 600 overlooked or cryptic species awaiting discovery through similar studies.

Table 2 Overlooked species of fishes found as a result of large barcoding studies in diverse ocean regions^a

Name of study	No. of species	No. of specimens	Overlooked species (N)	Overlooked species (%)	Reference
Fishes of Australia	207	754	5	2.40	Ward et al. 2005
Sharks and rays of Australia	210	945	4	1.90	Ward et al. 2008
Fishes of Pacific Canada	201	1,225	2	1.00	Steinke et al. 2009a
Reef-associated fishes	390	1,638	7	1.80	Steinke et al. 2009b
Arctic fishes of W. Pacific	106	882	3	2.80	Mecklenburg et al. 2010

^aThe studies listed examined more than 650 specimens of more than 100 species. Analysis resulted in detection of highly divergent barcodes within some putative species. Barcoding evidence of “overlooked” species was corroborated through morphological, geographical, and ecological data.

Reptiles. DNA barcodes can be useful for conservation and wildlife forensics of sea turtles by identifying turtle meat and eggs illegally traded or carcasses stranded on beaches (Vargas et al. 2009). Although sea turtles represent an ancient group with a slow mutation rate, all species were successfully identified and no cryptic species were revealed based on genetic distances and character-based methods (Naro-Maciel et al. 2010).

Birds. The CBOL All Birds Barcoding Initiative (<http://www.barcodingbirds.org>) includes seabirds. Among 17 sets of species of North American seabirds with overlapping barcodes, Kerr et al. (2007) found 8 species of large white-headed gulls. These may represent well-formed species that are losing genetic identity due to secondary contact and hybridization. The study also showed deep divergence between two lineages of the circum-Arctic species *Fulmaris glacialis*.

Mammals. Marine mammals currently lack a comprehensive library of DNA barcodes. Analysis of beaked whales (Ziphiidae), the least known of all cetacean families, led to significant discoveries, including description of a new species from the north Pacific, *Mesoplodon perrini* (Dalebout et al. 2002); resurrection of a long-forgotten species in the Southern Hemisphere, *M. traversii* (van Helden et al. 2002); confirmation of the identity of the enigmatic tropical bottlenose whale, *Indopacetus pacificus* (Dalebout et al. 2003); and perhaps discovery of a new species of *Mesoplodon* in the tropical Pacific (Dalebout et al. 2007).

Problem children. In two of the most ecologically important groups on coral reefs, the Anthozoa (e.g., corals and sea anemones) and Porifera (sponges), COI sequences have diverged too little to be diagnostic for all species (or even some genera) and hybrids cannot be distinguished from their maternal ancestors (Neigel et al. 2007). COI gene trees are typically paraphyletic (i.e., containing a single ancestor and some, but not all, of its descendents). Whereas in sponges a longer COI fragment than the standard 5' end might be useful (Erpenbeck et al. 2006), cnidarian barcoding might need another gene (Shearer & Coffroth 2008, Shearer et al. 2002).

Evolutionarily young (i.e., recently isolated) species present challenging cases for barcoding. Distinct but sympatric species may have very similar or identical barcodes due to incomplete lineage sorting or introgression (Kempainen et al. 2009, Steinke et al. 2009a). This was an early criticism of barcoding (Rubinoff et al. 2006), although such cases are rare outside the sponges, ctenophores, and anthozoans. Comparative analysis of a secondary nuclear marker can be used to confirm—or refute—the barcode results (e.g., Cardenas et al. 2009, Sonnenberg et al. 2007).

Two other groups bear brief mention here, although the difficulties may be more a matter of methodology than molecular evolution. Very few COI sequences are available for the pelagic

groups Appendicularia (Larvacea) and Thaliacea. Until sufficient COI sequence data are available and can be analyzed, it will remain unclear whether or not COI will be an appropriate barcode gene for these groups.

Barcoding Ocean Realms

Pelagic. Until recently, some pelagic taxa (e.g., copepods, euphausiids, and chaetognaths) were thought to be well known taxonomically. However, DNA barcoding of representatives of even these well-known groups has increasingly altered this perspective, and morphologically cryptic but genetically distinctive species of zooplankton are being found with increasing frequency. This issue is especially relevant for widely distributed species and/or for species with disjoint distributional ranges (McManus & Katz 2009). Interestingly, coastal and neritic environments appear to contain a higher frequency of zooplankton species that are endemic (i.e., occupying a highly restricted or unique habitat or geographic range) than do open ocean regions (e.g., Chen & Hare 2008, Dawson & Hamner 2005, Ueda & Bucklin 2006). DNA barcodes for pelagic assemblages have provided evidence of the power of COI for reliable identification of species across the 14 metazoan phyla and 20+ invertebrate classes occurring in the pelagic realm (Bucklin et al. 2010a,b) (**Figure 4**).

Polar seas. Polar regions are under threat from human activities, ocean acidification, and global warming, with potentially greater impacts than at lower latitudes. Arctic and Antarctic regions are considered to have intermediate levels of diversity (~5,000 species each), but these figures may markedly underestimate actual levels. Recent CoML and International Polar Year (IPY) sampling efforts have yielded new collections for DNA barcoding (Grant & Linse 2009, Mincks Hardy




Figure 4

Distance-based analysis of barcodes for a marine zooplankton assemblage of the Sargasso Sea, Northwest Atlantic Ocean. The tree shown was determined using the Neighbor Joining algorithm and Kimura-2-Parameter (K2P) genetic distances and was bootstrapped 1,000 times. A total of 328 individuals of 207 species were barcoded. The tree shows strong support for barcode resolution of species and major functional groups (usually classes) of zooplankton and fish. Figure modified from Bucklin et al. (2010b).

et al. 2010). The result is a growing collection of studies that employed barcodes both in the Arctic (Bucklin et al. 2010a, Mecklenburg et al. 2010) and Antarctic (Rock et al. 2008, Smith et al. 2008b, Webb et al. 2006). Increased sampling efforts in shallow and deep polar waters have uncovered novel species and generated numerous records of previously unrecorded taxa. Some of the new Arctic records may represent northward range expansions of boreal species with a warming climate, but many discoveries likely reflect the access to new samples from poorly sampled deeper areas. Using barcoding, Hunt et al. (2010) found species with bipolar distributions (i.e., occurring at both North and South polar regions) to be considerably more rare than previously proposed. Analysis revealed genetic structuring and cryptic speciation in bipolar species with both disjunct (i.e., absent at midlatitudes) and continuous distributions.

Coral reefs. Coral reefs are among the most endangered marine ecosystems, and dramatic declines in corals and fishes have been well documented (Gardner et al. 2003, Knowlton & Jackson 2008, Pandolfi et al. 2003). Most reef biodiversity lies outside these two groups in what is termed cryptofauna (i.e., hidden biodiversity associated with coral reefs), largely overlooked in biodiversity estimates. A study of the crustacean cryptofauna inhabiting dead coral heads of a single coral species revealed an unexpected high diversity of brachyuran crabs in a small geographic region (Plaisance et al. 2009). Similar findings have been reported for stomatopods (Barber & Boyce 2006) and even for fishes (Steinke et al. 2009b). Lins-de-Barros et al. (2010) used COI to confirm the presence of coral in a study of coral-associated archaea, bacteria, and eukaryotic algal plastid communities. The CoML coral reef project CREEFS has deployed Autonomous Reef Monitoring Structures (ARMS) in most of the world's coral seas and is developing DNA barcode catalogs for all of the species that settle on these structures.

Other marine communities. The Moorea Biocode Project is carrying out a comprehensive inventory of all nonmicrobial life of the French Polynesian island (Check 2006), using a barcode reference library as well as high-throughput sequencing for multigene analysis. Initial work on crustaceans uncovered a wealth of unrecognized diversity in the relatively well-known reef dweller *Calcinus* (Malay & Paulay 2010) and in crustacean cryptofauna (Plaisance et al. 2009). Hubert et al. (2010) created a reference collection of barcoded species of the perciform families Acanthuridae and Holocentridae; they found unprecedented levels of resolution in the identification of early stages of fish and provided evidence that conspecific larvae do not always aggregate in large schools.

BARCODE ANALYSIS BEYOND SPECIES IDENTIFICATION

Population Genetic and Phylogeographic Analysis

Population genetic analysis entails the quantitative description and statistical analysis of genetic diversity and structure (i.e., the distribution of genetic variation within and between populations) for a given species. Population genetic characters can also be used for phylogeographic analysis (i.e., the description of the geographical distributions of the genetic lineages within a population or species; Avise 2000). Whereas an ecologist might use archival or data-telemetry tags to track dispersal patterns, and a physical oceanographer might use drifters and dyes to analyze transport in currents, another useful approach to understanding dispersal is to use a tag that the organisms themselves carry—their COI barcode. Recent reviews of larval dispersal and population connectivity (Cowen & Sponaugle 2009) and gene flow (Hellberg 2009) in the ocean have provided comprehensive assessment and analyses for marine organisms. We note here a selection of studies

that have used the COI barcode region for population genetic and phylogeographic analysis of marine species. Indeed, COI sequence variation has proved useful to examine a wide variety of fundamental questions in the ecology and evolution of marine metazoa and has resolved geographic and temporal patterns over a range of scales

Gene flow and population connectivity. COI revealed significant population genetic structuring of marine metazoan species across the taxonomic spectrum, including sponges (Duran et al. 2004b, López-Legentil & Pawlik 2009), flatworms (Sanna et al. 2009), polychaetes (Breton et al. 2003, Xu et al. 2009), gastropods (Duda & Lee 2009, Gruenthal & Burton 2008, Keeney et al. 2009, Kelly & Eernisse 2007), bivalves (DeBoer et al. 2008, Lind et al. 2007), cephalopods (Teske et al. 2007a), crustaceans (Azuma et al. 2008, Liu et al. 2009, Luttikhuisen et al. 2008, Mathews 2007, Unal et al. 2006, Zitari-Chatte et al. 2009), brittle stars (Christensen et al. 2008, Muths et al. 2009), sea urchins (Duran et al. 2004a, Lessios et al. 2003), and ascidians (Pérez-Portela & Turon 2008a). Little or no differentiation for COI among geographic populations—even at global scales—was observed for a coralline sponge (Wörheide 2006), moon jelly (Ki et al. 2008), harpacticoid copepod (Eberl et al. 2007), commensal barnacle on hawksbill turtles (Torres-Pratts et al. 2009), spiny lobster (Inoue et al. 2007, Silberman et al. 1994), fiddler crab (Silva et al. 2010), galatheid shrimp (Samadi et al. 2006), amphidromous gastropod (Crandall et al. 2009), and mussel (Nicastro et al. 2008). Small- and large-scale patterns of connectivity are determined by many factors, including the ecology, life history, and behavior of the species. The rock-pool harpacticoid copepod *Tigriopus* has become a model system for small-scale population differentiation using COI (e.g., Handschumacher et al. 2010, Rawson & Burton 2006). For pelagic copepods, large-scale patterns of connectivity are strongly influenced by vertical migration behavior, which allows position-keeping and generates population differentiation (Blanco-Bercial et al. 2010) (Figure 5).

Historical range expansion and glacial refugia. COI is a frequent genetic marker of choice for phylogeographic and population genetic studies aimed at describing historical patterns of range expansion and contraction and at identifying glacial refugia for marine species. A number of North Atlantic marine species have been inferred to have shared patterns of preglacial maxima of demographic expansion, with evidence of glacial refugia and postglacial expansion (see Wares & Cunningham 2001). The European stalked barnacle, *Pollicipes pollicipes*, was found to exhibit this pattern based on COI variation (Campo et al. 2010). Similarly, the penaeid shrimp *Melicertus kerathurus* was hypothesized to have experienced a postglacial recolonization of the Mediterranean Sea from its Atlantic refugium based on COI variation (Pellerito et al. 2009). Species restricted to the nearshore and coastal margins of the North Atlantic have been hypothesized to exhibit marked cladogenesis in COI resulting from range expansion from glacial refugia, including the copepod *Acartia tonsa* (Hill 2004) and the amphipod *Gammarus tigrinus* (Kelly et al. 2006).

Van den Broeck et al. (2008) proposed a stepping stone postglacial range expansion through the Eastern Atlantic to explain marked phylogeographic structuring of COI of the periwinkle *Tectarius striatus*. Elsewhere, Goodall-Copestake et al. (2010) timed population expansion of Southern Ocean krill based on COI variation. Coral Triangle seastars and associated parasitic and commensal species revealed species-specific phylogeographic patterns resulting from differential postglacial range expansions (Crandall et al. 2008). Species-specific phylogeographic patterns for intertidal gastropod (Littorinidae) species throughout the Indo-West Pacific were tied to historical events (Reid et al. 2006).




Figure 5

Connectivity of two calanoid copepod species populations in the Atlantic Ocean based on the cytochrome oxidase subunit I (COI) barcode region. *Metridia lucens* (left) shows lower connectivity; the species' strong vertical migration behavior may aid geographic position-keeping and reduce large-scale dispersal; nonmigrating *Nannocalanus minor* (right) shows greater connectivity throughout the ocean basins; it may be more readily transported in surface currents (Blanco-Bercial et al. 2010). Haplotype network diagrams were determined by TCS, a computer program that estimates gene genealogies (Clement et al. 2000). Colors indicate the geographic source of collections; sizes of pie diagrams and ovals are relative to numbers of haplotypes; dotted lines indicate numbers of nucleotide substitutions between haplotypes. Adapted from unpublished data and figure from L. Blanco-Bercial and A. Bucklin, Univ. of Conn.

Phylogeny

In general, the COI barcode region is a poor choice as a phylogenetic marker. However, on its own, the phylogenetic signal of COI has been widely shown to be useful at two disparate levels of analysis: resolution of relationships among very closely related taxa and resolution of nodes between major groups of animals. Browne et al. (2007) sampled a wide range of amphipods and presented a preliminary molecular phylogeny based on COI; many lineage relationships between taxa within three hyperiid clades were highly supported. COI resolved relationships among giant clam species, which were shown to be monophyletic by Nuryanto et al. (2007). COI resolved deeper phylogenetic relationships among corals (Fukami et al. 2004, 2008).

It has been widely suggested that the inclusion of a large number of genomic regions in phylogenetic analysis—in a total evidence approach—will improve the quality of the inferences (Cummings et al. 1995). However, different markers—even different mitochondrial genes—differ markedly in amounts of nucleotide variation and rates of divergence, thus generating significant incongruence in the resultant tree topologies. The advantages of combining data partitions despite failed homogeneity tests are debatable (e.g., Giribet 2002); lack of congruence with other mitochondrial and nuclear makers has not prevented publication of innumerable studies that include COI in multigene phylogenetic analyses, frequently also including large- (28S) and small-subunit (18S) rRNA sequences.

There are examples of both congruence and lack of congruence of COI with other genes for molecular phylogenetic analysis. Compared with a reference topology for neogastropods based on whole mitochondrial genomes, Cunha et al. (2009) found that COI showed the lowest number of congruent internal branches and thus the lowest phylogenetic performance. Combined analysis of 28S rRNA and COI yielded highly congruent patterns of relationship at highly supported nodes within the sponge class Demospongia (Nichols 2005). Cárdenas et al. (2010) used COI to resolve phylogenetic relationships of sponges of the family Geodiidae and found congruent results with other gene regions, including 28S rRNA. COI provided a phylogenetic signal (identified in the combined analysis with 18S and 28S rRNA) for resolution of relationships within the isopod family Munnopsidae (Osborn 2009). Except for the resolution of basal taxa, Meland & Willassen (2004) found similar tree topologies for 18S rRNA and COI for species of the deep-sea mysid genus *Pseudomma*.

Trophic Relationships (Gut Contents)

Confirmation that short barcodes are diagnostic of species (Hajibabaei et al. 2006b) paved the way for their use in circumstances where full barcodes are impossible to obtain, as for degraded DNA in predator gut contents. Protocols are now available for barcoding in circumstances that necessitate dealing with fragmented DNA such as gut content and food web analysis (Carreon-Martinez & Heath 2010). Töbe et al. (2010) developed oligonucleotide probes based on barcodes for dominant copepods to detect the species in the guts of the larval krill *Euphausia superba* collected from the Lazarev Sea in the Southern Ocean; real-time, quantitative PCR results showed that the krill were feeding exclusively on *Oithona* sp. Krill was also the subject of an earlier study that employed short sequences of small-subunit (18S) rRNA to identify diatom prey from adult animals in the Indian Ocean (Passmore et al. 2006). In order to directly measure predation rates by older copepod stages upon copepod nauplii, Durbin et al. (2008) used species-specific primers and quantitative PCR to calculate ingestion rates based on DNA disappearance rates. The benthic marine worm *Xenoturbella* was found to be frequently contaminated with molluscan DNA; barcoding data (Bourlat et al. 2008) showed that the molluscan DNA originated from bivalves living in the same environment, indicating that the worm fed specifically on bivalve prey.

Dunn et al. (2010) conducted a diet study of deepwater sharks utilizing full length barcodes to identify prey species; they estimated that between 84 and 223 stomachs were needed to measure 90% of the extrapolated total prey richness. DNA identification was used to characterize prey species of large pelagic fishes by Smith et al. (2005). Especially for predators with diverse diets, group-specific primers, usually based on more conserved genes than COI, can expand the range of dietary items detected in gut contents. As an example, marine vertebrate diets were examined by Jarman et al. (2004) using probes based on 28S rRNA.

Forensics and Seafood Safety

DNA barcoding can be used for cooked or processed seafood. Smith et al. (2008a) obtained species-level identifications for smoked fish fillets of several genera; seafood sampled at commercial markets and restaurants was mislabeled at least in 25% of the cases (Barbuto et al. 2010, Cohen et al. 2009, Lowenstein et al. 2009, Marko et al. 2004, Wong & Hanner 2008). COI barcoding has been validated as a diagnostic marker for species identification (Dawnay et al. 2007) and is under consideration by the Food and Drug Administration in the United States as a replacement for the technique of protein isoelectric focusing for fish and fish product identification (Cohen et al. 2009, Yancy et al. 2007).

In one of the first applications of DNA barcoding in a human health context, barcodes were used to identify tuna sushi to the species level concomitant with mercury testing (Lowenstein et al. 2009). DNA barcodes enabled researchers to determine which species warranted inclusion in consumer advisories or trade restrictions, and whether the data used by health agencies accurately reflected mercury levels.

Marine Invasive Species

Barcodes can provide accurate identification of marine invasive species. In some instances, COI variation can be analyzed to reveal the geographic source region and pathway of invasion. Genetic approaches to the study of marine invasions were reviewed recently by Geller et al. (2009), who summarize barcoding studies as well. Two studies that showcase the usefulness of COI in this regard include a study of invasive pathways of a bryozoan (Mackie et al. 2006) and comparative analysis of COI diversity of native and introduced populations of an amphipod (Ashton et al. 2008).

METHODOLOGY—PRESENT AND FUTURE

High-Throughput Sequencing and Environmental Barcoding

Barcoding is moving to massively parallel sequencing platforms to speed analysis of COI from mixed genomes, as well as whole genomes for the target species. Barcodes for larval fishes have been successfully determined using high-throughput sequencing (Richardson et al. 2006). High-throughput sequencing is also useful for determining COI barcodes from bulk environmental samples. Sometimes called environmental barcoding, this methodology offers an accurate, cost-effective, and rapid estimation of species diversity of marine communities based on barcodes. DNA or mRNA is extracted from bulk environmental samples and analyzed for COI sequence diversity. High-throughput sequencing for exhaustive analysis of cDNA libraries can help reduce artifacts due to mispriming and pseudogenes. Also, short (<100 bp) sequences are sufficient to identify species in most cases by matching to a comprehensive taxon- or region-specific database of DNA barcodes of known species. As increasingly complete DNA barcode databases become available, increasingly accurate estimation of new or undescribed species will be possible. Machida et al. (2009) demonstrated the power of environmental barcoding by analyzing a zooplankton sample collected in the Equatorial Pacific Ocean; they detected 189 species among 828 COI sequences, using threshold differences of 9% between species and distance-based clustering. Creer et al. (2010) analyzed the highly diverse but generally cryptic meiofaunal assemblage using environmental barcoding.

Real-Time Quantitative PCR

A natural extension of DNA barcoding is the use of the barcode sequence to design probes for real-time, or quantitative, PCR (QPCR), which can be used to estimate absolute and relative abundances of species in unsorted bulk samples. Pan et al. (2008) used real-time PCR assays based on COI for detection and relative quantification of decapod crustacean larvae.

DNA Microarrays

DNA microarrays are matrices with a detector DNA sequence (probe) affixed in each cell, which will bind (hybridize) to the labeled target DNA in the test sample. DNA barcodes can be used to design probe sequences for up to ~7,000 species, although new analytical software may be

required. The main limitation of the array-based approach is that it requires advance knowledge of sequences in target species. Because of a lack of exact matches, undiscovered haplotypes or geographic variants could fail to anneal properly to the probes on the array. This problem could be avoided by providing a set of several different probes per species, which limits the use of microarrays for large-scale biodiversity projects. However, DNA microarrays made for regional assemblages have the potential to accelerate routine at-sea or lab-based taxonomic analysis of marine samples for known species.

The *in silico* study on the effectiveness of mammalian COI and cytochrome *b* (*cyt b*) sequences for probe design suggested that both genes yield a high number of probes (Hajibabaei et al. 2007a). However, as the behavior of oligonucleotide probes in hybridisation experiments cannot be predicted, *in silico* results should be viewed with caution. Another study suggests that COI and *cyt b* are not well suited for probe design in fish species (Kochzius et al. 2010), and similar findings were reported for COI in fungi (Chen et al. 2009). Sequences of mitochondrial 12S and 16S rRNA may be more suitable for the design of functional probes in fish species (Kochzius et al. 2008). This result is particularly interesting as COI is markedly superior for sequence-based species identification of fishes; 16S rRNA frequently fails to discriminate closely related species. It is likely that microarray-based identification of marine metazoan species will require probes designed not solely from COI but from a combination of genes.

Sources of Error

Heteroplasmy. Strict maternal inheritance of mtDNA should result in homoplasmy (i.e., only one mtDNA type within a cell or individual). Heteroplasmy (i.e., the presence of a mixture of more than one type of mtDNA within a cell or individual) can be caused by the differential amplification of nuclear pseudogenes (Bensasson et al. 2001), paternal leakage (Kvist et al. 2003), somatic mutation (Moum & Bakke 2001), and cross-species transfer (Barbara et al. 2007). Mussels of the *Mytilus edulis* species group appear to be exceptions to usual mtDNA inheritance patterns; homoplasmic individuals with mtDNA similar to the heteroplasmic mtDNA types have been found (Hoeh et al. 1991). These observations are best explained by the hypothesis that biparental inheritance of mitochondrial DNA can occur in *Mytilus* (e.g., Ort & Pogson 2007, Śmietanka et al. 2009). Further complexity in interpreting *Mytilus* barcodes and other mtDNA patterns is introduced by recombination after hybridization between species (see Riginos & Henzler 2008).

Pseudogenes. Nuclear mitochondrial pseudogenes, or numts (i.e., nonfunctional copies of mtDNA in the nucleus), have been found in major clades of eukaryotic organisms. Studies have revealed high frequencies in some taxa (e.g., the snapping shrimp *Alpheus*; Williams & Knowlton 2001). They can be easily coamplified with orthologous mtDNA when using conserved universal primers. This is problematical for DNA barcoding, particularly as nucleotide substitutions are not expected to occur as rapidly in the nucleus as in the mitochondrial genome (Brown et al. 1979). However, recent studies (Smith et al. 2007, Song et al. 2008) have presented strategies to reduce the risk of misinterpreting barcoding results.

Selection. Although the barcode region is frequently interpreted as a selectively neutral marker of species identity, numerous studies have confirmed the influence of selection on patterns of diversity and rates of divergence (e.g., Meiklejohn et al. 2007). A marine barcoding study by Foltz et al. (2004) found less COI diversity among species of the star fish *Leptasterias* that brood their young, indicating the influence of life history factors on patterns of COI nucleotide substitutions. Galtier et al. (2009) reviewed evidence for mitochondrial selective sweeps (i.e., reduction or elimination of

sequence variation resulting from recent and strong natural selection) and hypothesized that levels of intraspecific mtDNA divergence may primarily reflect the time elapsed since the last selective sweep. Selective sweeps would thus interfere with interpretations of demographic processes based on mtDNA but could aid species identification using barcodes by reducing intraspecific variation and making the barcoding gap more pronounced.

The formalin problem. Barcoding of marine organisms has been slowed by preservation and storage of samples in formalin solutions, which are not always adequately buffered for pH. Both the aqueous matrix and the rapid acidification of unbuffered formalin have exacerbated the difficulty of recovering DNA from archived samples (Tang 2006). Long-term storage in buffered formalin does not preclude some types of molecular genetic analysis, assuming caution is paid to possible sources of artifact (Zimmermann et al. 2008). Numerous protocols exist to recover DNA from formalin-preserved tissue; barcodes obtained from formalin-preserved marine specimens have been reported for zooplankton (Bucklin & Allen 2004, Kirby & Lindley 2005) and fish (Zhang 2010). Even if new protocols can be developed for DNA recovery from formalin-preserved tissue, marine specimens intended for molecular analysis should be preserved and maintained in ethanol or be deep frozen (Tang 2006).

Buyer beware. The rapid expansion of COI sequences in the NCBI GenBank database has provided an increasingly useful and comprehensive library for species identification of diverse marine metazoans. However, closer examination can reveal numerous errors and artifacts with various causes, including taxonomic misidentification of species, DNA sequencing quality control concerns (e.g., sequencing or manual editing errors, stop codons, and insertions and deletions in reading frames), unrecognized pseudogenes, and microbial contaminants (e.g., Siddall et al. 2009), among others. Errors are becoming more evident as barcode databases are more densely populated. Erroneous sequences can be identified by comparative examination within a particular taxonomic group; for example, Buhay (2009) reported numerous errors after examination of crustacean COI sequences on GenBank.

FUTURE ISSUES FOR DNA BARCODING

1. The new taxonomy. Concern that only a tiny cadre of expert taxonomists can identify and describe marine species has become the basis for renewed public interest and reaffirmed understanding of the scientific importance of species-level biodiversity throughout the global ocean. We can expect DNA barcoding to continue to rebuild the field of taxonomy and retain—or recover—the body of knowledge of marine biodiversity.
2. The Rosetta Stone for marine biodiversity. The growing database of gold standard DNA barcodes serves as a Rosetta Stone for marine biodiversity, providing the keys to link species names, morphology, and DNA sequence variation. The COI sequence library will serve to translate or decipher the complex array of morphological characters that are used to describe and discriminate species by traditional taxonomists (Hajibabaei et al. 2007b, Stoeckle & Hebert 2008).
3. Environmental barcoding. Analysis of marine biodiversity will proceed through barcoding of specimens that have not been—or cannot be—identified by recognized taxonomic authorities. Some specimens may be new or undescribed species,

whereas others may be simply damaged or poorly preserved. In many cases, high-throughput sequencing will profoundly outstrip the speed with which traditional sample analysis can be performed. Barcodes from unidentified specimens are being used to analyze species-level diversity and estimate numbers of novel species. New analytical approaches may eventually allow accurate classification of unknown barcodes.

4. Whole-genome barcoding. We may expect that, in the not-too-distant-future, it may become both possible and cost-effective to characterize whole genomes (or whole mitochondrial genomes) for the same purposes now served by the COI barcode region for species identification, discrimination, and discovery.
5. Conservation of biodiversity. Until recently, the prospects for rapidly cataloging tens of thousands species associated with hot spot habitats were very dim. Amid genuine fears that species extinction will outpace discovery, DNA barcoding offers the promise of greatly accelerating the rate of biodiversity analysis for threatened or endangered marine habitats with exceptionally high species diversity, such as coral reefs (Neigel et al. 2007).
6. Ocean monitoring and observing. O'Dor et al. (2010) summarized a comprehensive analysis of new technologies for ocean observing, including the use of DNA barcoding for rapid and remote analysis of species diversity of ocean ecosystems. Operational approaches to marine biodiversity that use barcodes to monitor ecosystem health, inform fisheries management, and detect species diversity for near real-time ocean observing are on the horizon.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

We gratefully acknowledge the programmatic support of the Alfred P. Sloan Foundation. Invaluable support and assistance for the effort described here was provided by the Census of Marine Life and the Consortium for the Barcode of Life.

LITERATURE CITED

- Abdo Z, Golding B. 2007. A step toward barcoding life: a model based, decision theoretic method to assign genes to pre-existing species groups. *Syst. Biol.* 56:44–56
- Allcock AL, Barratt I, Eleaume M, Linse K, Norman MD, et al. 2010. Cryptic speciation and the circumpolarity debate: a case study on endemic Southern Ocean octopuses using the COI barcode of life. *Deep-Sea Res. II*. In review
- Altschul SF, Gish W, Miller W, Meyers EW, Lipman DJ. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215:403–10
- Ashton GV, Stevens MI, Hart MC, Green DH, Burrows MT, et al. 2008. Mitochondrial DNA reveals multiple Northern Hemisphere introductions of *Caprella mutica* (Crustacea, Amphipoda). *Mol. Ecol.* 17:1293–303
- Avise JC. 2000. *Phylogeography: The History and Formation of Species*. Cambridge, MA: Harvard Univ. Press. 447 pp.
- Avise JC. 2004. *Molecular Markers, Natural History and Evolution*. Sunderland, MA: Sinauer. 541 pp.

- Azuma N, Kunihiro Y, Sasaki J, Mihara E, Mihara Y, et al. 2008. Genetic variation and population structure of hair crab (*Erimacrus isenbeckii*) in Japan inferred from mitochondrial DNA sequence analysis. *Mar. Biotechnol.* 10:39–48
- Barbara T, Palma-Silva C, Paggi GM, Bered F, Fay MF, et al. 2007. Cross-species transfer of nuclear microsatellite markers: potential and limitations. *Mol. Ecol.* 16:3759–67
- Barber P, Boyce SL. 2006. Estimating diversity of Indo-Pacific coral reef stomatopods through DNA barcoding of stomatopod larvae. *Proc. R. Soc. Lond. B* 273:2053–61
- Barbuto M, Galimberti A, Ferri E, Labra M, Malandra R, et al. 2010. DNA barcoding reveals fraudulent substitutions in shark seafood products: the Italian case of “palombo” (*Mustelus* spp.). *Food Res. Int.* 43:376–81
- Barroso R, Klautau M, Solé-Cava A, Paiva P. 2010. *Eurythoe complanata* (Polychaeta: Amphinomidae), the ‘cosmopolitan’ fireworm, consists of at least three cryptic species. *Mar. Biol.* 157:69–80
- Bensasson D, Zhang D, Hartl D, Hewitt G. 2001. Mitochondrial pseudogenes: evolution’s misplaced witnesses. *Trends Ecol. Evol.* 16:314–21
- Bertolazzi P, Felici G, Weitschek E. 2009. Learning to classify species with barcodes. *BMC Bioinformatics* 10:S7
- Bhadury P, Austen MC. 2010. Barcoding marine nematodes: an improved set of nematode 18S rRNA primers to overcome eukaryotic cointerference. *Hydrobiologia* 641:245–51
- Bhadury P, Austen MC, Bilton DT, Lamshead PJD, Rogers AD, et al. 2006. Development and evaluation of a DNA-barcoding approach for the rapid identification of nematodes. *Mar. Ecol. Prog. Ser.* 320:1–9
- Blanco-Bercial L, Bucklin A, Viñas MD, Cornils A. 2010. Zooplankton phylogeography as a measure of Atlantic Ocean ecosystem connectivity. Paper presented at *Ocean Sciences Meeting*, Portland, Ore
- Blanquer A, Uriz M-J. 2007. Cryptic speciation in marine sponges evidenced by mitochondrial and nuclear genes: A phylogenetic approach. *Mol. Phylog. Evol.* 45:392–97
- Blaxter M, Mann J, Chapman T, Thomas F, Whitton C, et al. 2005. Defining operational taxonomic units using DNA barcode data. *Philos. Trans. R. Soc. Lond. B* 360:1935–43
- Böttger-Schnack R, Machida R. 2010. Comparison of morphological and molecular traits for species identification and taxonomic grouping of oncaeid copepods. *Hydrobiologia* 2010:1–15
- Bouchet P. 2006. The magnitude of marine biodiversity. In *The exploration of Marine Biodiversity: Scientific and Technical Challenges*, ed. CM Duarte, pp. 31–64. Fundacion BBVA, Madrid
- Bourlat SJ, Nakano H, Akerman M, Telford MJ, Thorndyke MC, et al. 2008. Feeding ecology of *Xenoturbella bocki* (phylum Xenoturbellida) revealed by genetic barcoding. *Mol. Ecol. Resour.* 8:18–22
- Breton S, Dufresne F, Desrosiers G, Blier PU. 2003. Population structure of two northern hemisphere polychaetes, *Neanthes virens* and *Hediste diversicolor* (Nereididae), with different life-history traits. *Mar. Biol.* 142:707–15
- Brown WM, George MJr, Wilson AC. 1979. Rapid evolution of animal mitochondrial DNA. *Proc. Natl. Acad. Sci. USA* 76:1967–71
- Browne WE, Haddock SHD, Martindale MQ. 2007. Phylogenetic analysis of lineage relationships among hyperiid amphipods as revealed by examination of the mitochondrial gene, cytochrome oxidase I (COI). *Integr. Comp. Biol.* 47:815–30
- Bucklin A, Allen LD. 2004. MtDNA sequencing from zooplankton after long-term preservation in buffered formalin. *Mol. Phylog. Evol.* 30:879–82
- Bucklin A, Frost BW. 2009. Morphological and molecular phylogenetic analysis of evolutionary lineages within *Clausocalanus* (Copepoda: Calanoida). *J. Crust. Biol.* 29:111–20
- Bucklin A, Frost BW, Bradford-Grieve J, Allen LD, Copley NJ. 2003. Molecular systematic and phylogenetic assesment of 34 calanoid copepod species of the Calanidae and Clausocalanidae. *Mar. Biol.* 142:333–43
- Bucklin A, Hopcroft RR, Kosobokova KN, Nigro LM, Ortman BD, et al. 2010a. DNA barcoding of Arctic Ocean holozooplankton for species identification and recognition. *Deep-Sea Res. II* 57:40–48
- Bucklin A, Ortman BD, Jennings RM, Nigro L, Sweetman CJ, et al. 2010b. A “Rosetta Stone” for zooplankton: DNA barcode analysis of holozooplankton diversity of the Sargasso Sea (NW Atlantic Ocean). *Deep-Sea Res. II*. In press; doi: 10.1016/j.dsr2.2010.09.025
- Bucklin A, Wiebe PH, Smolenack SB, Copley NJ, Beaudet JG, et al. 2007. DNA barcodes for species identification of euphausiids (Euphausiacea, Crustacea). *J. Plankton Res.* 29:483–93

- Buhay JE. 2009. "COI-Like" sequences are becoming problematic in molecular systematic and DNA barcoding studies. *J. Crust. Biol.* 29:96–110
- Calderón I, Garrabou J, Aurelle D. 2006. Evaluation of the utility of COI and ITS markers as tools for population genetic studies of temperate gorgonians. *J. Exp. Mar. Biol. Ecol.* 336:184–97
- Campo D, Molares J, Garcia L, Fernandez-Rueda P, Garcia-Gonzalez C, et al. 2010. Phylogeography of the European stalked barnacle (*Pollicipes pollicipes*): identification of glacial refugia. *Mar. Biol.* 157:147–56
- Cárdenas P, Menegola C, Rapp HT, Diaz MC. 2009. Morphological description and DNA barcodes of shallow-water *Tetractinellida* (Porifera: Demospongiae) from Bocas del Toro, Panama, with description of a new species. *Zootaxa* 2009:1–39
- Cárdenas P, Rapp HT, Schander C, Tendal OS. 2010. Molecular taxonomy and phylogeny of the Geodiidae (Porifera, Demospongiae, Astrophorida)—Combining phylogenetic and Linnaean classification. *Zool. Scr.* 39:89–106
- Carreon-Martinez L, Heath DD. 2010. Revolution in food web analysis and trophic ecology: diet analysis by DNA and stable isotope analysis. *Mol. Ecol.* 19:25–27
- Carstensen D, Laudien J, Leese F, Arntz W, Held C. 2009. Genetic variability, shell and sperm morphology suggest that the surf clams *Donax marincovichii* and *D. obesulus* are one species. *J. Mollus. Stud.* 75:381–90
- Check E. 2006. Treasure island: pinning down a model ecosystem. *Nature* 439:378–79
- Chen G, Hare MP. 2008. Cryptic ecological diversification of a planktonic estuarine copepod, *Acartia tonsa*. *Mol. Ecol.* 17:1451–68
- Chen W, Seifert KA, Levesque CA. 2009. A high density COX1 barcode oligonucleotide array for identification and detection of species of *Penicillium* subgenus *Penicillium*. *Mol. Ecol. Resour.* 9:114–29
- Christensen A, Christensen E, Weisrock D. 2008. Population genetic structure of North American *Ophiactis* spp. brittle stars possessing hemoglobin. *Mar. Biol.* 154:755–63
- Chu KH, Xu M, Li CP. 2009. Rapid DNA barcoding analysis of large datasets using the composition vector method. *BMC Bioinformatics* 10:S8
- Clare E, Kerr K, von Königslöw T, Wilson J, Hebert P. 2008. Diagnosing mitochondrial DNA diversity: applications of a sentinel gene approach. *J. Mol. Evol.* 66:362–67
- Clement M, Posada D, Crandall K. 2000. TCS: a computer program to estimate gene genealogies. *Mol. Ecol.* 9:1657–60
- Cohen NJ, Deeds JR, Wong ES, Hanner R, Yancy HF, et al. 2009. Public health response to puffer fish (Tetrodotoxin) poisoning from mislabeled product. *J. Food Prot.* 72:810–17
- Costa FO, Henzler CM, Lunt DH, Whiteley NM, Rock J. 2009. Probing marine *Gammarus* (Amphipoda) taxonomy with DNA barcodes. *Syst. Biodivers.* 7:365–79
- Cowen RK, Sponaugle S. 2009. Larval dispersal and marine population connectivity. *Annu. Rev. Marine Sci.* 1:443–66
- Crandall ED, Jones ME, Muñoz MM, Akinronbi B, Erdmann MV, et al. 2008. Comparative phylogeography of two seastars and their ectosymbionts within the Coral Triangle. *Mol. Ecol.* 17:5276–90
- Crandall ED, Taffel JR, Barber PH. 2009. High gene flow due to pelagic larval dispersal among South Pacific archipelagos in two amphidromous gastropods (Neritimorpha: Neritidae). *Heredity* 104:563–74
- Creer S, Fonseca VG, Porazinska DL, Giblin-Davis RM, Sung W, et al. 2010. Ultrasequencing of the meiofaunal biosphere: practice, pitfalls and promises. *Mol. Ecol.* 19:4–20
- Cross M, Collins C, Campbell N, Watts P, Chubb J, et al. 2007. Levels of intrahost and temporal sequence variation in a large CO 1 subunits from *Anisakis simplex* sensu stricto (Rudolphi 1809) (Nematoda: Anisakidae): implications for fisheries management. *Mar. Biol.* 151:695–702
- Cummings M, Otto S, Wakeley J. 1995. Sampling properties of DNA sequence data in phylogenetic analysis. *Mol. Biol. Evol.* 12:814–22
- Cunha R, Grande C, Zardoya R. 2009. Neogastropod phylogenetic relationships based on entire mitochondrial genomes. *BMC Evol. Biol.* 9:210
- D'Amato M, Harkins G, de Oliveira T, Teske P, Gibbons M. 2008. Molecular dating and biogeography of the neritic krill *Nyctiphanes*. *Mar. Biol.* 155:243–47
- Dalebout ML, Baker CS, Anderson RC, Best PB, Cockcroft VG, et al. 2003. Appearance, distribution, and genetic distinctiveness of longman's beaked whale, *Indopacetus pacificus*. *Mar. Mamm. Sci.* 19:421–61

- Dalebout ML, Baker CS, Steel D, Robertson KM, Chivers SJ, et al. 2007. A divergent mtDNA lineage among *Mesoplodon* beaked whales: Molecular evidence for a new species in the Tropical Pacific? *Mar. Mamm. Sci.* 23:954–66
- Dalebout ML, Mead JG, Baker CS, Baker AN, Helden AL. 2002. A new species of beaked whale *Mesoplodon perrini* sp. n. (Cetacea: Ziphiidae) discovered through phylogenetic analyses of mitochondrial DNA sequences. *Mar. Mamm. Sci.* 18:577–608
- Dawnay N, Ogden R, McEwing R, Carvalho GR, Thorpe RS. 2007. Validation of the barcoding gene CO1 for use in forensic genetic species identification. *Forensic Sci. Int.* 173:1–6
- Dawson MN. 2005a. *Cyanea capillata* is not a cosmopolitan jellyfish: morphological and molecular evidence for *C. annaskala* and *C. rosea* (Scyphozoa: Semaeostomeae: Cyaneidae) in south-eastern Australia. *Invertebr. Syst.* 19:361–70
- Dawson MN. 2005b. Morphologic and molecular redescription of *Catostylus mosaicus conservativus* (Scyphozoa: Rhizostomeae: Catostylidae) from south-east Australia. *J. Mar. Biol. Assoc. U.K.* 85:723–31
- Dawson MN. 2005c. Renaissance taxonomy: integrative evolutionary analyses in the classification of Scyphozoa. *J. Mar. Biol. Assoc. U.K.* 85:733–39
- Dawson MN, Hamner WM. 2005. Rapid evolutionary radiation of marine zooplankton in peripheral environments. *Proc. Natl. Acad. Sci. USA* 102:9235–40
- Dawson MN, Jacobs DK. 2001. Molecular evidence for cryptic species of *Aurelia aurita* (Cnidaria, Scyphozoa). *Biol. Bull.* 200:92–96
- DeBoer TS, Subia MD, Erdmann MV, Kovitvoingsa K, Barber PH. 2008. Phylogeography and limited genetic connectivity in the endangered boring giant clam across the coral triangle. *Conserv. Biol.* 22:1255–66
- Derycke S, Ley PD, Ley ITD, Holovachov O, Rigaux A, et al. 2010. Linking DNA sequences to morphology: cryptic diversity and population genetic structure in the marine nematode *Thoracostoma trachygaster* (Nematoda, Leptosomatidae). *Zool. Scr.* 39:276–89
- DeSalle R. 2005. Conservation genetics—Genetics at the brink of extinction. *Heredity* 94:386–87
- DeSalle R, Egan MG, Siddall M. 2005. The unholy trinity: taxonomy, species delimitation and DNA barcoding. *Philos. Trans. R. Soc. Lond. B* 360:1905–16
- DeSalle ROB. 2006. Species discovery versus species identification in DNA barcoding efforts: response to Rubinoff. *Conserv. Biol.* 20:1545–47
- Dippenaar SM, Mathibela RB, Bloomer P. 2010. Cytochrome oxidase I sequences reveal possible cryptic diversity in the cosmopolitan symbiotic copepod *Nesippus orientalis* Heller, 1868 (Pandaridae: Siphonostomatoida) on elasmobranch hosts from the KwaZulu-Natal coast of South Africa. *Exp. Parasitol.* 125:42–50
- Duda T, Lee T. 2009. Isolation and population divergence of a widespread Indo-West Pacific marine gastropod at Easter Island. *Mar. Biol.* 156:1193–202
- Dunn MR, Szabo A, McVeagh MS, Smith PJ. 2010. The diet of deepwater sharks and the benefits of using DNA identification of prey. *Deep-Sea Res. I.* 57:923–30
- Duran S, Palacin C, Becerro MA, Turon X, Giribet G. 2004a. Genetic diversity and population structure of the commercially harvested sea urchin *Paracentrotus lividus* (Echinodermata, Echinoidea). *Mol. Ecol.* 13:3317–28
- Duran S, Pascual M, Turon X. 2004b. Low levels of genetic variation in mtDNA sequences over the western Mediterranean and Atlantic range of the sponge *Crambe crambe* (Poecilosclerida). *Mar. Biol.* 144:31–35
- Durbin E, Casas M, Rynearson T, Smith D. 2008. Measurement of copepod predation on nauplii using qPCR of the cytochrome oxidase I gene. *Mar. Biol.* 153:699–707
- Eberl R, Cohen S, Cipriano F, Carpenter EJ. 2007. Genetic diversity of the pelagic harpacticoid copepod *Macrosetella gracilis* on colonies of the cyanobacterium *Trichodesmium* spp. *Aquat. Biol.* 1:33–43
- Eddy SR. 1998. Profile hidden Markov models. *Bioinformatics* 14:755–63
- Elsasser SC, Floyd R, Hebert PDN, Schulte-Hostedde AI. 2009. Species identification of North American guinea worms (Nematoda: Draconulcus) with DNA barcoding. *Mol. Ecol. Resour.* 9:707–12
- Erpenbeck D, Duran S, Rützler K, Paul V, Hooper JNA, et al. 2007. Towards a DNA taxonomy of Caribbean demosponges: a gene tree reconstructed from partial mitochondrial CO1 gene sequences supports previous rDNA phylogenies and provides a new perspective on the systematics of Demospongiae. *J. Mar. Biol. Assoc. U.K.* 87:1563–70

- Erpenbeck D, Hooper JNA, Worheide G. 2006. COI phylogenies in diploblasts and the 'Barcoding of Life'—Are we sequencing a suboptimal partition? *Mol. Ecol. Notes* 6:550–53
- Erséus C, Kvist S. 2007. COI variation in Scandinavian marine species of *Tubificoides* (Annelida: Clitellata: Tubificidae). *J. Mar. Biol. Assoc. U.K.* 87:1121–26
- Feng Y, Li Q, Kong L, Zheng X. 2010. DNA barcoding and phylogenetic analysis of Pectinidae (Mollusca: Bivalvia) based on mitochondrial COI and 16S rRNA genes. *Mol. Biol. Rep.* In press; doi:10.1007/s11033-010-0107-1
- Ferguson JWH. 2002. On the use of genetic divergence for identifying species. *Biol. J. Linn. Soc.* 75:509–16
- Ferri E, Barbuto M, Bain O, Galimberti A, Uni S, et al. 2009. Integrated taxonomy: traditional approach and DNA barcoding for the identification of filarioid worms and related parasites (Nematoda). *Front. Zool.* 6:1
- Floyd RM, Rogers AD, Lamshead PJD, Smith CR. 2005. Nematode-specific PCR primers for the 18S small subunit rRNA gene. *Mol. Ecol. Notes* 5:611–12
- Folino-Rorem N, Darling J, D'Ausilio C. 2009. Genetic analysis reveals multiple cryptic invasive species of the hydrozoan genus *Cordylophora*. *Biol. Invasions* 11:1869–82
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol. Mar. Biol. Biotechnol.* 3:294–99
- Foltz DW, Hrnicevich AW, Rocha-Olivares A. 2004. Apparent selection intensity for the cytochrome oxidase subunit I gene varies with mode of reproduction in echinoderms. *Genetica* 122:115–25
- Fukami H, Budd AF, Paulay G, Solé-Cava A, Allen Chen C, Iwao K, Knowlton N. 2004. Conventional taxonomy obscures deep divergence between Pacific and Atlantic corals. *Nature* 427:832–35
- Fukami H, Chen CA, Budd AF, Collins A, Wallace C, Chuang YY, et al. 2008. Mitochondrial and nuclear genes suggest that stony corals are monophyletic but most families of stony corals are not (Order Scleractinia, Class Anthozoa, Phylum Cnidaria). *PLoS One* 3:e3222
- Galtier N, Nabholz B, Glémin S, Hurst GDD. 2009. Mitochondrial DNA as a marker of molecular diversity: a reappraisal. *Mol. Ecol.* 18:4541–50
- Gardner TA, Cote IM, Gill JA, Grant A, Watkinson AR. 2003. Long-term region-wide declines in Caribbean corals. *Science* 301:958–60
- Geller JB, Darling JA, Carlton JT. 2009. Genetic perspectives on marine biological invasions. *Annu. Rev. Marine Sci.* 2:367–93
- Giribet G. 2002. Current advances in the phylogenetic reconstruction of metazoan evolution. A new paradigm for the Cambrian explosion? *Mol. Phylog. Evol.* 24:345–57
- Gomez A, Hughes RN, Wright PJ, Carvalho GR, Lunt DH. 2007a. Mitochondrial DNA phylogeography and mating compatibility reveal marked genetic structuring and speciation in the NE Atlantic bryozoan *Celleporella hyalina*. *Mol. Ecol.* 16:2173–88
- Gomez A, Wright PJ, Lunt DH, Cancino JM, Carvalho GR, et al. 2007b. Mating trials validate the use of DNA barcoding to reveal cryptic speciation of a marine bryozoan taxon. *Proc. R. Soc. Lond. B* 274:199–207
- Goodall-Copetake WP, Perez-Espona S, Clark MS, Murphy EJ, Seear PJ, et al. 2010. Swarms of diversity at the gene *cox1* in Antarctic krill. *Heredity* 104:513–18
- Govindarajan AF, Boero F, Halanych KM. 2006. Phylogenetic analysis with multiple markers indicates repeated loss of the adult medusa stage in Campanulariidae (Hydrozoa, Cnidaria). *Mol. Phylog. Evol.* 38:820–34
- Grant RA, Linse K. 2009. Barcoding Antarctic biodiversity: current status and the CAML initiative, a case study of marine invertebrates. *Polar Biol.* 32:1629–37
- Gruenthal KM, Burton RS. 2008. Genetic structure of natural populations of the California black abalone (*Haliotis cracherodii* Leach, 1814), a candidate for endangered species status. *J. Exp. Mar. Biol. Ecol.* 355:47–58
- Hajibabaei M, Janzen DH, Burns JM, Hallwachs W, Hebert PDN. 2006a. DNA barcodes distinguish species of tropical Lepidoptera. *Proc. Natl. Acad. Sci. USA* 103:968–71
- Hajibabaei M, Singer GAC, Clare EL, Hebert PDN. 2007a. Design and applicability of DNA arrays and DNA barcodes in biodiversity monitoring. *BMC Biol.* 5:24
- Hajibabaei M, Singer GAC, Hebert PD, Hickey DA. 2007b. DNA barcoding: how it complements taxonomy, molecular phylogenetics and population genetics. *Trends Genet.* 23:167–72

- Hajibabaei M, Smith MA, Janzen DH, Rodriguez JJ, Whitfield JB, et al. 2006b. A minimalist barcode can identify a specimen whose DNA is degraded. *Mol. Ecol. Notes* 6:959–64
- Handschumacher L, Steinarsdóttir M, Edmands S, Ingólfsson A. 2010. Phylogeography of the rock-pool copepod *Tigriopus brevicornis* (Harpacticoida) in the northern North Atlantic, and its relationship to other species of the genus. *Mar. Biol.* 157:1357–66
- Havermans C, Nagy ZT, Sonet G, De Broyer C, Martin P. 2010. Incongruence between molecular phylogeny and morphological classification in amphipod crustaceans: a case study of Antarctic lysianassoids. *Mol. Phylog. Evol.* 55:202–09
- Hebert PDN, Cywinska A, Ball SL, deWaard JR. 2003a. Biological identifications through DNA barcodes. *Proc. R. Soc. Lond. B* 270:313–21
- Hebert PDN, Penton EH, Burns JM, Janzen DH, Hallwachs W. 2004a. Ten species in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astraptes fulgerator*. *Proc. Natl. Acad. Sci. USA* 101:14812–17
- Hebert PDN, Ratnasingham S, deWaard JR. 2003b. Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proc. R. Soc. Lond. B* 270:S96–99
- Hebert PDN, Stoeckle MY, Zemplak TS, Francis CM. 2004b. Identification of birds through DNA barcodes. *PLoS Biol.* 2:e312
- Heim I, Nickel M, Brümmer F. 2007. Molecular markers for species discrimination in poriferans: a case study on species of the genus *Aplysina*. In *Série Livros*, ed. MR Custódio, G Lôbo-Hajdu, E Hajdu, G Muricy. Museu Nacional, Rio de Janeiro
- Helgen LE, Rouse GW. 2006. Species delimitation and distribution in *Aporometra* (Crinoidea: Echinodermata): endemic Australian featherstars. *Invertebr. Syst.* 20:395–414
- Hellberg M. 2006. No variation and low synonymous substitution rates in coral mtDNA despite high nuclear variation. *BMC Evol. Biol.* 6:24
- Hellberg ME. 2009. Gene flow and isolation among populations of marine animals. *Annu. Rev. Ecol. Evol. Syst.* 40:291–310
- Hickerson MJ, Meyer CP, Moritz C. 2006. DNA Barcoding will often fail to discover new animal species over broad parameter space. *Syst. Biol.* 55:729–39
- Hill RS. 2004. *Genetic diversity and structure of calanoid copepods: molecular evolutionary patterns in coastal estuaries (Acartia tonsa) and the open ocean (Calanus spp.)*. Univ. of New Hampshire, Durham, NH. 143 pp.
- Hill RS, Allen LD, Bucklin A. 2001. Multiplexed species-specific PCR protocol to discriminate four N. Atlantic *Calanus* species, with a mtCOI gene tree for ten *Calanus* species. *Mar. Biol.* 139:279–87
- Hirose M, Hirose E. 2009. DNA barcoding in photosymbiotic species of *Diplosoma* (Ascidiacea: Didemnidae), with the description of a new species from the Southern Ryukyus, Japan. *Zool. Sci.* 26:564–68
- Hirose M, Osawa M, Hirose E. 2010. DNA barcoding of hermit crabs of genus *Clibanarius* Dana, 1852 (Anomura: Diogenidae) in the Ryukyu Islands, southwestern Japan. *Zootaxa*: 59–66
- Hoareau TB, Boissin E. 2010. Design of phylum-specific hybrid primers for DNA barcoding: addressing the need for efficient COI amplification in the Echinodermata. *Mol. Ecol. Resour.* In press; doi: 10.1111/j.1755-0998.2010.02848.x
- Hoeh WR, Blakley KH, Brown WM. 1991. Heteroplasmy suggests limited biparental inheritance of *Mytilus* mitochondrial DNA. *Science* 251:1488–90
- Holland BS, Dawson MN, Crow GL, Hofmann DK. 2004. Global phylogeography of *Casiopea* (Scyphozoa: Rhizostomeae): molecular evidence for cryptic species and multiple invasions of the Hawaiian Islands. *Mar. Biol.* 145:1119–28
- Holmes BH, Steinke D, Ward RD. 2009. Identification of shark and ray fins using DNA barcoding. *Fisheries Res.* 95:280–88
- Hou Z-E, Li Z, Li S-q. 2009. Identifying Chinese species of *Gammarus* (Crustacea: Amphipoda) using DNA barcoding. *Curr. Zool.* 55:158–64
- Hou Z, Fu JH, Li SQ. 2007. A molecular phylogeny of the genus *Gammarus* (Crustacea: Amphipoda) based on mitochondrial and nuclear gene sequences. *Mol. Phylog. Evol.* 45:596–611
- Huang D, Meier R, Todd PA, Chou LM. 2008. Slow mitochondrial COI sequence evolution at the base of the metazoan tree and its implications for DNA barcoding. *J. Mol. Evol.* 66:167–74

- Hubert N, Delrieu-Trottin E, Irisson J-O, Meyer C, Planes S. 2010. Identifying coral reef fish larvae through DNA barcoding: a test case with the families Acanthuridae and Holocentridae. *Mol. Phylogen. Evol.* 55:1195–203
- Hunt B, Strugnell J, Bednarsek N, Linse K, Nelson RJ, et al. 2010. Poles Apart: The “bipolar” pteropod species *Limacina helicina* is genetically distinct between the Arctic and Antarctic oceans. *PLoS ONE* 5:e9835
- Inoue N, Watanabe H, Kojima S, Sekiguchi H. 2007. Population structure of Japanese spiny lobster *Panulirus japonicus* inferred by nucleotide sequence analysis of mitochondrial COI gene. *Fish. Sci.* 73:550–56
- Ivanova NV, Dewaard JR, Hebert PDN. 2006. An inexpensive, automation-friendly protocol for recovering high-quality DNA. *Mol. Ecol. Notes* 6:998–1002
- Ivanova NV, Zemlak TS, Hanner R, Hebert PDN. 2007. Universal primer cocktails for fish DNA barcoding. *Mol. Ecol. Notes* 7:544–48
- Jarman SN, Deagle BE, Gales NJ. 2004. Group-specific polymerase chain reaction for DNA-based analysis of species diversity and identity in dietary samples. *Mol. Ecol.* 13:1313–22
- Jennings RM, Bucklin A, Ossenbrügger H, Hopcroft RR. 2010a. Analysis of genetic diversity of planktonic gastropods from several ocean regions using DNA barcodes. *Deep-Sea Res. II*. In press
- Jennings RM, Bucklin A, Pierrot-Bults A. 2010b. Barcoding of arrow worms (phylum Chaetognatha) from three oceans: genetic diversity and evolution within an enigmatic phylum. *PLoS ONE* 5:e9949
- Johnson SB, Warén A, Vrijenhoek RC. 2008. DNA Barcoding of *Lepetodrilus* limpets reveals cryptic species. *J. Shellfish Res.* 27:43–51
- Jones WJ, Macpherson E. 2007. Molecular phylogeny of the East Pacific squat lobsters of the genus *Munidopsis* (Decapoda: Galatheidae) with the descriptions of seven new species. *J. Crust. Biol.* 27:477–501
- Keeney DB, King TM, Rowe DL, Poulin R. 2009. Contrasting mtDNA diversity and population structure in a direct-developing marine gastropod and its trematode parasites. *Mol. Ecol.* 18:4591–603
- Kelly DW, MacIsaac HJ, Heath DD. 2006. Vicariance and dispersal effects on phylogeographic structure and speciation in a widespread estuarine invertebrate. *Evolution* 60:257–67
- Kelly RP, Eernisse DJ. 2007. Southern hospitality: a latitudinal gradient in gene flow in the marine environment. *Evolution* 61:700–7
- Kelly RP, Sarkar IN, Eernisse DJ, DeSalle ROB. 2007. DNA barcoding using chitons (genus *Mopalia*). *Mol. Ecol. Notes* 7:177–83
- Kemppainen P, Panova M, Hollander J, Johannesson K. 2009. Complete lack of mitochondrial divergence between two species of NE Atlantic marine intertidal gastropods. *J. Evol. Biol.* 22:2000–11
- Kerr K, Stoeckle M, Dove C, Weigt L, Francis C, et al. 2007. Comprehensive DNA barcode coverage of North American birds. *Mol. Ecol. Notes* 7:535–43
- Ki J-S, Hwang D-S, Shin K, Yoon WD, Lim D, et al. 2008. Recent moon jelly (*Aurelia* sp.1) blooms in Korean coastal waters suggest global expansion: examples inferred from mitochondrial COI and nuclear ITS-5.8S rDNA sequences. *ICES J. Mar. Sci.* 65:443–52
- Kirby RR, Lindley JA. 2005. Molecular analysis of Continuous Plankton Recorder samples, an examination of echinoderm larvae in the North Sea. *J. Mar. Biol. Assoc. U.K.* 85:451–59
- Knowlton N, Jackson JBC. 2008. Shifting baselines, local impacts, and global change on coral reefs. *PLoS Biol.* 6:e54
- Kochzius M, Nölte M, Weber H, Silkenbeumer N, Hjørleifsdóttir S, et al. 2008. DNA microarrays for identifying fishes. *Mar. Biotechnol.* 10:207–17
- Kochzius M, Seidel C, Antoniou A, Kumar Botla S, Campo D, et al. 2010. Identifying fishes through DNA barcodes and microarrays. *PLoS ONE*. In review
- Koski LB, Golding B. 2001. The closest BLAST hit is often not the nearest neighbor. *J. Mol. Ecol.* 52:540–42
- Krabbe K, Leese F, Mayer C, Tollrian R, Held C. 2010. Cryptic mitochondrial lineages in the widespread pycnogonid *Colossendeis megalonyx* Hoek, 1881 from Antarctic and Subantarctic waters. *Polar Biol.* 33:281–92
- Krebes L, Blank M, Jurss K, Zettler ML, Bastrop R. 2010. Glacial-driven vicariance in the amphipod *Gammarus duebeni*. *Mol. Phylogen. Evol.* 54:372–85
- Krug PJ, Ellingson RA, Burton R, Valdes A. 2007. A new *Poecilognous* species of sea slug (Opisthobranchia: Sacoglossa) from California: comparison with the planktotrophic congener *Alderia modesta*. *J. Molluscan Stud.* 73:29–38

- Kvist L, Martens J, Nazarenko AA, Orell M. 2003. Paternal leakage of mitochondrial DNA in the great tit (*Parus major*). *Mol. Biol. Evol.* 20:243–47
- Lambshhead PJD. 2001. Marine nematode diversity. In *Nematology, Advances and Perspectives*, ed. ZX Chen, SY Chen, DW Dickson. Wallingford, UK: CABI Publ.
- Last PR, Gledhill DC, Holmes BH. 2007a. A new handfish, *Brachionichthys australis* sp. nov. (Lophiiformes: Brachionichthyidae), with a redescription of the critically endangered spotted handfish, *B. hirsutus* (Lacepede). *Zootaxa* 1666:53–68
- Last PR, White WT, Pogonoski JJ. 2007b. *Descriptions of New Dogfishes of the Genus Squalus (Squaloidea: Squalidae)*. CSIRO, Hobart, Tasmania
- Lessios HA, Kane J, Robertson DR. 2003. Phylogeography of the pantropical sea urchin *Tripneustes*: contrasting patterns of population structure between oceans. *Evolution* 57:2026–36
- Lind CE, Evans BS, Taylor JJU, Jerry DR. 2007. Population genetics of a marine bivalve, *Pinctada maxima*, throughout the Indo-Australian Archipelago shows differentiation and decreased diversity at range limits. *Mol. Ecol.* 16:5193–203
- Lins-de-Barros MM, Vieira RP, Cardoso AM, Monteiro VA, Turque AS, et al. 2010. Archaea, Bacteria, and algal plastids associated with the reef-building corals *Siderastrea stellata* and *Mussismilia hispida* from Buzios, South Atlantic Ocean, Brazil. *Microb. Ecol.* 59:523–32
- Liu Y, Liu R, Ye L, Liang J, Xuan F, et al. 2009. Genetic differentiation between populations of swimming crab *Portunus trituberculatus* along the coastal waters of the East China Sea. *Hydrobiologia* 618:125–37
- López-Legentil S, Pawlik J. 2009. Genetic structure of the Caribbean giant barrel sponge *Xestospongia muta* using the I3-M11 partition of COI. *Coral Reefs* 28:157–65
- Lorion J, Duperron Sb, Gros O, Cruaud C, Samadi S. 2009. Several deep-sea mussels and their associated symbionts are able to live both on wood and on whale falls. *Proc. R. Soc. Lond. B* 276:177–85
- Lowenstein JH, Amato G, Kolokotronis SO. 2009. The real *maccoyii*: identifying tuna sushi with DNA barcodes—contrasting characteristics attributes and genetic distances. *PLoS ONE* 4(11):e7866
- Luttikhuisen PC, Campos J, Bleijswijk Jv, Peijnenburg KTCA, van der Veer HW. 2008. Phylogeography of the common shrimp, *Crangon crangon* (L.) across its distribution range. *Mol. Phylog. Evol.* 46:1015–30
- Luttikhuisen PC, Dekker R. 2010. Pseudo-cryptic species *Arenicola defodiens* and *Arenicola marina* (Polychaeta: Arenicolidae) in Wadden Sea, North Sea and Skagerrak: morphological and molecular variation. *J. Sea Res.* 63:17–23
- Machida R, Hashiguchi Y, Nishida M, Nishida S. 2009. Zooplankton diversity analysis through single-gene sequencing of a community sample. *BMC Genomics* 10:438
- Machida R, Miya M, Nishida M, Nishida S. 2006. Molecular phylogeny and evolution of the pelagic copepod genus *Neocalanus* (Crustacea: Copepoda). *Mar. Biol.* 148:1071–79
- Mackie J, Keough M, Christidis L. 2006. Invasion patterns inferred from cytochrome oxidase I sequences in three bryozoans, *Bugula neritina*, *Watersipora subtorquata*, and *Watersipora arcuata*. *Mar. Biol.* 149:285–95
- Malay MCD, Paulay G. 2010. Peripatric speciation drives diversification and distributional pattern of reef hermit crabs (Decapoda: Diogenidae: *Calcinus*). *Evolution* 64:634–62
- Marko PB, Lee SC, Rice AM, Gramling JM, Fitzhenry TM, McAlister JS, et al. 2004. Mislabeling of a depleted reef fish. *Nature* 430:309–10
- Mathews L. 2007. Evidence for restricted gene flow over small spatial scales in a marine snapping shrimp *Alpheus angulosus*. *Mar. Biol.* 152:645–55
- Matz MV, Nielsen R. 2005. A likelihood ratio test for species membership based on DNA sequence data. *Philos. Trans. R. Soc. Lond. B* 360:1969–74
- McManus GB, Katz LA. 2009. Molecular and morphological methods for identifying plankton: What makes a successful marriage? *J. Plankton Res.* 31:1119–29
- Mecklenburg CW, Möller PR, Steinke D. 2010. Biodiversity of arctic marine fishes: taxonomy and zoogeography. *Mar. Biodivers.* In review
- Meier R, Shiyang K, Vaidya G, Ng PK. 2006. DNA barcoding and taxonomy in Diptera: a tale of high intraspecific variability and low identification success. *Syst. Biol.* 55:715–28
- Meiklejohn CD, Montooth KL, Rand DM. 2007. Positive and negative selection on the mitochondrial genome. *Trends Genet.* 23:259–63

- Meland K, Willassen EE. 2004. Molecular phylogeny and biogeography of the genus *Pseudomma* (Peracarida: Mysida). *J. Crust. Biol.* 24:541–57
- Meyer CP, Paulay G. 2005. DNA barcoding: Error rates based on comprehensive sampling. *PLoS Biol.* 3:2229–38
- Mikkelsen NT, Schander C, Willassen E. 2007. Local scale DNA barcoding of bivalves (Mollusca): a case study. *Zool. Scr.* 36:455–63
- Miller SE. 2007. DNA barcoding and the renaissance of taxonomy. *Proc. Natl. Acad. Sci. USA* 104:4775–76
- Min XJ, Hickey DA. 2007. Assessing the effect of varying sequence length on DNA barcoding of fungi. *Mol. Ecol. Notes* 7:365–73
- Mincks Hardy S, Carr CM, Hardman M, Steinke D, Corstorphine E, et al. 2010. Biodiversity and phylogeography of Arctic marine fauna: insights from molecular tools. *Mar. Biodivers.* In review
- Miyamoto H, Machida R, Nishida S. 2010. Genetic diversity and cryptic speciation of the deep sea chaetognath *Caecosagitta macrocephala*. *Deep-Sea Res. II*. In press
- Moritz C, Cicero C. 2004. DNA barcoding: promise and pitfalls. *PLoS Biol.* 2:1529–31
- Moszczyńska A, Locke SA, McLaughlin JD, Marcogliese DJ, Crease TJ. 2009. Development of primers for the mitochondrial cytochrome c oxidase I gene in digenetic trematodes (Platyhelminthes) illustrates the challenge of barcoding parasitic helminths. *Mol. Ecol. Resour.* 9:75–82
- Moum T, Bakke I. 2001. Mitochondrial control region structure and single site heteroplasmy in the razorbill (*Alca torda*; Aves). *Curr. Genet.* 39:198–203
- Moura CJ, Harris DJ, Cunha MR, Rogers AD. 2008. DNA barcoding reveals cryptic diversity in marine hydroids (Cnidaria, Hydrozoa) from coastal and deep-sea environments. *Zool. Scr.* 31:93–108
- Muths D, Jollivet D, Gentil F, Davoult D. 2009. Large-scale genetic patchiness among NE Atlantic populations of the brittle star *Ophiothrix fragilis*. *Aquat. Biol.* 5:117–32
- Naro-Maciel E, Reid B, Fitzsimmons NN, Le M, Desalle R, et al. 2010. DNA barcodes for globally threatened marine turtles: a registry approach to documenting biodiversity. *Mol. Ecol. Resour.* 10:252–63
- Naughton KM, O'Hara TD. 2009. A new brooding species of the biscuit star *Tosia* (Echinodermata: Asteroidea: Goniasteridae), distinguished by molecular, morphological and larval characters. *Invertebr. Syst.* 23:348–66
- Neigel J, Domingo A, Stake J. 2007. DNA barcoding as a tool for coral reef conservation. *Coral Reefs* 26:487–99
- Nicastro K, Zardi G, McQuaid C, Teske P, Barker N. 2008. Coastal topography drives genetic structure in marine mussels. *Mar. Ecol. Prog. Ser.* 368:189–95
- Nichols SA. 2005. An evaluation of support for order-level monophyly and interrelationships within the class Demospongiae using partial data from the large subunit rDNA and cytochrome oxidase subunit I. *Mol. Phylog. Evol.* 34:81–96
- Nielsen R, Matz M. 2006. Statistical approaches for DNA barcoding. *Syst. Biol.* 55:162–69
- Nuryanto A, Duryadi D, Soedharma D, Blohm D. 2007. Molecular phylogeny of giant clams based on mitochondrial DNA Cytochrome C Oxidase I gene. *HAYATI* 14:162–66
- O'Dor R, Acosta J, Bergstad OA, Brainard R, Bratley J, et al. 2010. Bringing life to Ocean Observation. In *Proceedings of OceanObs'09: Sustained Ocean Observations and Information for Society*, ed. J Hall, DE Harrison, D Stammer, vol. 1. Ecol. Soc. Am., Ithaca, NY
- Øines Ø, Heuch PA. 2005. Identification of sea louse species of the genus *Caligus* using mtDNA. *J. Mar. Biol. Assoc. U.K.* 85:73–79
- Ort BS, Pogson GH. 2007. Molecular population genetics of the male and female mitochondrial DNA molecules of the California sea mussel, *Mytilus californianus*. *Genetics* 177:1087–99
- Ortman BD, Bucklin A, Pagés F, Youngbluth M. 2010. DNA barcoding of the Medusozoa. *Deep-Sea Res. II*. In press
- Osborn KJ. 2009. Relationships within the Munnopsidae (Crustacea, Isopoda, Asellota) based on three genes. *Zool. Scr.* 38:617–35
- Paine MA, McDowell JR, Graves JE. 2008. Specific identification using COI sequence analysis of scombrid larvae collected off the Kona coast of Hawaii Island. *Ichthyol. Res.* 55:7–16
- Paine MA, McDowell JR, Graves JE. 2007. Specific identification of Western Atlantic Ocean scombrids using mitochondrial DNA cytochrome c oxidase subunit I (COI) gene region sequences. *Bull. Mar. Sci.* 80:353–67

- Pan M, McBeath A, Hay S, Pierce G, Cunningham C. 2008. Real-time PCR assay for detection and relative quantification of *Liocarcinus depurator* larvae from plankton samples. *Mar. Biol.* 153:859–70
- Pandolfi JM, Bradbury RH, Sala E, Hughes TP, Bjorndal KA, et al. 2003. Global trajectories of the long-term decline of coral reef ecosystems. *Science* 301:955–58
- Pardo LM, Ampuero D, Véliz D. 2009. Using morphological and molecular tools to identify megalopae larvae collected in the field: the case of sympatric *Cancer* crabs. *J. Mar. Biol. Assoc. U.K.* 89:481–90
- Park MH, Sim CJ, Baek J, Min GS. 2007. Identification of genes suitable for DNA barcoding of morphologically indistinguishable Korean Halichondriidae sponges. *Mol. Cells* 23:220–27
- Passmore AJ, Jarman SN, Swadling KM, Kawaguchi S, McMinn A, et al. 2006. DNA as a dietary biomarker in Antarctic krill, *Euphausia superba*. *Mar. Biotechnol.* 8:686–96
- Pegg GG, Sinclair B, Briskey L, Aspden WJ. 2006. MtDNA barcode identification of fish larvae in the southern Great Barrier Reef, Australia. *Sci. Mar.* 70:7–12
- Pellerito R, Arculeo M, Bonhomme F. 2009. Recent expansion of Northeast Atlantic and Mediterranean populations of *Melicertus (Penaeus) kerathurus* (Crustacea: Decapoda). *Fish. Sci.* 75:1089–95
- Pérez-Portela R, Bishop JDD, Davis AR, Turon X. 2009. Phylogeny of the families Pyuridae and Styelidae (Stolidobranchiata, Ascidiacea) inferred from mitochondrial and nuclear DNA sequences. *Mol. Phylog. Evol.* 50:560–70
- Pérez-Portela R, Duran S, Palaćin C, Turon X. 2007. The genus *Pycnoclavella* (Ascidiacea) in the Atlanto-Mediterranean region: a combined molecular and morphological approach. *Invertebr. Syst.* 21:187–205
- Pérez-Portela R, Turon X. 2008a. Cryptic divergence and strong population structure in the colonial invertebrate *Pycnoclavella communis* (Ascidiacea) inferred from molecular data. *Zoology* 111:163–78
- Pérez-Portela R, Turon X. 2008b. Phylogenetic relationships of the Clavelinidae and Pycnoclavellidae (Ascidiacea) inferred from mtDNA data. *Invertebr. Biol.* 127:108–20
- Plaisance L, Knowlton N, Paulay G, Meyer C. 2009. Reef-associated crustacean fauna: biodiversity estimates using semiquantitative sampling and DNA barcoding. *Coral Reefs* 28:977–86
- Pöppe J, Sutcliffe P, Hooper JN, Wörheide G, Erpenbeck D. 2010. CO I barcoding reveals new clades and radiation patterns of Indo-Pacific sponges of the family Irciniidae (Demospongiae: Dictyoceratida). *PLoS ONE* 5:e9950
- Puillandre N, Maylac M, Boisselier M-C, Cruaud C, Samadi S. 2009a. An integrative approach to species delimitation in *Benthomangelia* (Mollusca: Conoidea). *Biol. J. Linn. Soc.* 96:696–708
- Puillandre N, Samadi S, Boisselier M-C, Cruaud C, Bouchet P. 2009b. Molecular data provide new insights on the phylogeny of the Conoidea (Neogastropoda). *Nautilus* 123:202–10
- Puillandre N, Strong EE, Bouchet P, Boisselier M-C, Couloux A, et al. 2009c. Identifying gastropod spawn from DNA barcodes: possible but not yet practicable. *Mol. Ecol. Resour.* 9:1311–21
- Pyle RL, Earle JL, Greene BD. 2008. Five new species of the damselfish genus *Chromis* (Perciformes: Labroidae: Pomacentridae) from deep coral reefs in the tropical western Pacific. *Zootaxa* 1671:3–31
- Rach J, DeSalle R, Sarkar IN, Schierwater B, Hadrys H. 2008. Character-based DNA barcoding allows discrimination of genera, species and populations in Odonata. *Philos. Trans. R. Soc. Lond. B* 275:237–47
- Radulovici AE, Sainte-Marie B, Dufresne F. 2009. DNA barcoding of marine crustaceans from the Estuary and Gulf of St Lawrence: a regional-scale approach. *Mol. Ecol. Res.* 9:181–187
- Raleigh J, Redmond NE, Delahan E, Torpey S, van Soest RWM, et al. 2007. Mitochondrial cytochrome oxidase 1 phylogeny supports alternative taxonomic scheme for the marine Haplosclerida. *J. Mar. Biol. Assoc. U.K.* 87:1577–84
- Ratnasingham S, Hebert PDN. 2007. The barcode of life database. *Mol. Ecol. Notes* 7:355–64
- Raupach MJ, Mayer C, Malyutina M, Waegle JW. 2009. Multiple origins of deep-sea Asellota (Crustacea: Isopoda) from shallow waters revealed by molecular data. *Proc. R. Soc. Lond. B* 276:799–808
- Rawson PD, Burton RS. 2006. Molecular adaptation at the cytochrome c oxidase subunit 2 gene among divergent populations of the intertidal copepod, *Tigriopus californicus*. *J. Molec. Evol.* 62:753–64
- Reid DG, Lal K, Mackenzie-Dodds J, Kaligis F, Littlewood DTJ, et al. 2006. Comparative phylogeography and species boundaries in *Echinolittorina* snails in the central Indo-West Pacific. *J. Biogeogr.* 33:990–1006
- Remerie T, Bourgeois T, Peelaers D, Vierstraete A, Vanfleteren J, et al. 2006. Phylogeographic patterns of the mysid *Mesopodopsis slabberi* (Crustacea, Mysida) in Western Europe: evidence for high molecular diversity and cryptic speciation. *Mar. Biol.* 149:465–81

- Rice SA, Karl S, Rice KA. 2008. The *Polydora cornuta* complex (Annelida: Polychaeta) contains populations that are reproductively isolated and genetically distinct. *Invertebr. Biol.* 127:45–64
- Richardson D, Van Wye J, Exum A, Cowen R, Crawford D. 2006. High-throughput species identification: from DNA isolation to bioinformatics. *Mol. Ecol. Notes* 7:199–207
- Riginos C, Henzler C. 2008. Patterns of mtDNA diversity in North Atlantic populations of the mussel *Mytilus edulis*. *Mar. Biol.* 155:399–412
- Rock J, Costa FO, Walker DI, North AW, Hutchinson WF, et al. 2008. DNA barcodes of fish of the Scotia Sea, Antarctica indicate priority groups for taxonomic and systematics focus. *Antarct. Sci.* 20:253–62
- Ross HA, Murugan S, Sibon Li WL. 2008. Testing the reliability of genetic methods of species identification via simulation. *Syst. Biol.* 57:216–30
- Rubinoff D, Cameron S, Will K. 2006. A genomic perspective on the shortcomings of mitochondrial DNA for “barcoding” identification. *J. Hered.* 97:581–94
- Samadi S, Bottan L, Macpherson E, De Forges B, Boisselier M-C. 2006. Seamount endemism questioned by the geographic distribution and population genetic structure of marine invertebrates. *Mar. Biol.* 149:1463–75
- Sanna D, Lai T, Francalacci P, Curini-Galletti M, Casu M. 2009. Population structure of the *Monocelis lineata* (Proseriata, Monocelididae) species complex assessed by phylogenetic analysis of the mitochondrial Cytochrome c Oxidase subunit I (COI) gene. *Genet. Mol. Biol.* 32:864–67
- Sarkar IN, Planet PJ, Bael TE, Stanley SE, Siddall M, et al. 2002a. Characteristic attributes in cancer microarrays. *J. Biomed. Inf.* 35:111–22
- Sarkar IN, Planet PJ, DeSalle R. 2008. CAOS software for use in character-based DNA barcoding. *Mol. Ecol. Resour.* 8:1256–59
- Sarkar IN, Thornton J, Planet PJ, Schierwater B, DeSalle R. 2002b. A systematic method for classification of novel homeoboxes. *Mol. Phylogen. Evol.* 24:388–99
- Savolainen V, Cowan RS, Vogler AP, Roderick GK, Lane R. 2005. Towards writing the encyclopedia of life: an introduction to DNA barcoding. *Philos. Trans. R. Soc. Lond. B* 360:1805–11
- Shearer TL, Coffroth MA. 2008. Barcoding corals: limited by interspecific divergence, not intraspecific variation. *Mol. Ecol. Resour.* 8:247–55
- Shearer TL, Oppen MJHv, Romano SL, Wörheide G. 2002. Slow mitochondrial DNA sequence evolution in the Anthozoa (Cnidaria). *Mol. Ecol.* 11:2475–87
- Shih HT, Kamrani E, Davie PJF, Liu MY. 2009. Genetic evidence for the recognition of two fiddler crabs, *Uca iranica* and *U. albimana* (Crustacea: Brachyura: Ocypodidae), from the northwestern Indian Ocean, with notes on the *U. lactea* species-complex. *Hydrobiologia* 635:373–82
- Siddall ME, Fontanella FM, Watson SC, Kvist S, Erseus C. 2009. Barcoding bamboozled by bacteria: convergence to metazoan mitochondrial primer targets by marine microbes. *Syst. Biol.* 58:445–51
- Silberman JD, Sarver SK, Walsh PJ. 1994. Mitochondrial DNA variation and population structure in the spiny lobster *Panulirus argus*. *Mar. Biol.* 120:601–8
- Silva I, Mesquita N, Paula J. 2010. Lack of population structure in the fiddler crab *Uca annulipes* along an East African latitudinal gradient: genetic and morphometric evidence. *Mar. Biol.* 157:1113–26
- Sinniger F, Reimer JD, Pawlowski J. 2010. The Parazoanthidae (Hexacorallia: Zoantharia) DNA taxonomy: description of two new genera. *Mar. Biodivers.* 40:57–70
- Sinniger F, Reimer JD, Pawlowski J. 2008. Potential of DNA sequences to identify zoanthids (Cnidaria: Zoantharia). *Zool. Sci.* 25:1253–60
- Sirovich L, Stoeckle MY, Zhang Y. 2009. A scalable method for analysis and display of DNA sequences. *PLoS ONE* 4:e7051
- Sirovich L, Stoeckle MY, Zhang Y. 2010. Structural analysis of biodiversity. *PLoS ONE* 5:e9266
- Śmietanka B, Burzyński A, Wenne R. 2009. Molecular population genetics of male and female mitochondrial genomes in European mussels *Mytilus*. *Mar. Biol.* 156:913–25
- Smith M, Wood D, Janzen D, Hallwachs W, Hebert P. 2007. DNA barcodes affirm that 16 species of apparently generalist tropical parasitoid flies (Diptera, Tachinidae) are not all generalists. *Proc. Natl. Acad. Sci. USA* 104:4967–72
- Smith PJ, McVeagh MS, Steinke D. 2008a. Application of DNA barcoding for the identification of smoked fish products. *J. Fish Biol.* 72:464–71

- Smith PJ, McVeagh SM, Allain V, Sanchez C. 2005. DNA identification of gut contents of large pelagic fishes. *J. Fish Biol.* 67:1178–83
- Smith PJ, Steinke D, McVeagh MS, Stewart AL, Struthers CD, et al. 2008b. Molecular analysis of Southern Ocean skates (*Bathyraja*) reveals a new species of Antarctic skate. *J. Fish Biol.* 73:1170–82
- Song H, Buhay JE, Whiting MF, Crandall KA. 2008. Many species in one: DNA barcoding overestimates the number of species when nuclear mitochondrial pseudogenes are coamplified. *Proc. Natl. Acad. Sci. USA* 105:13486–491
- Sonnenberg R, Nolte A, Tautz D. 2007. An evaluation of LSU rDNA D1–D2 sequences for their use in species identification. *Front. Zool.* 4:6
- Spencer HG, Waters JM, Eichhorst TE. 2007. Taxonomy and nomenclature of black nerites (Gastropoda: Neritimorpha: *Nerita*) from the South Pacific. *Invertebr. Syst.* 21:229–37
- Spies IB, Gaichas S, Stevenson DE, Orr JW, Canino MF. 2006. DNA-based identification of Alaska skates (*Amblyraja*, *Bathyraja* and *Raja*: Rajidae) using cytochrome c oxidase subunit I (COI) variation. *J. Fish. Biol.* 69:283–92
- Steinke D, Vences M, Salzburger W, Meyer A. 2005. TaxI: a software tool for DNA barcoding using distance methods. *Philos. Trans. R. Soc. Lond. B* 360:1975–80
- Steinke D, Zemlak TS, Boutillier JA, Hebert PDN. 2009a. DNA barcoding Pacific Canada's fishes. *Mar. Biol.* 156:2641–47
- Steinke D, Zemlak TS, Hebert PDN. 2009b. Barcoding Nemo: DNA-based identifications for the ornamental fish trade. *PLoS ONE* 4:e6300
- Stoeckle MY, Hebert PDN. 2008. Barcode of Life: DNA Tags Help Classify Animals. *Sci. Am.* 299:82
- Sundberg P, Vodoti ET, Strand M. 2010. DNA barcoding should accompany taxonomy—the case of *Cerebratulus* spp (Nemertea). *Mol. Ecol. Resour.* 10:274–81
- Tang EPY. 2006. *Path to Effective Recovering of DNA from Formalin-Fixed Biological Samples in Natural History Collections: Workshop Summary*. Washington, DC: Natl. Acad. Press
- Tang RWK, Yau C, NG W-C. 2010. Identification of stomatopod larvae (Crustacea: Stomatopoda) from Hong Kong waters using DNA barcodes. *Mol. Ecol. Resour.* 10:439–48
- Teske PR, Oosthuizen A, Papadopoulos I, Barker N. 2007a. Phylogeographic structure of *Octopus vulgaris* in South Africa revisited: identification of a second lineage near Durban harbor. *Mar. Biol.* 151:2119–22
- Teske PR, Barker NP, McQuaid CD. 2007b. Lack of genetic differentiation among four sympatric southeast african intertidal limpets (Siphonariidae): phenotypic plasticity in a single species? *J. Molluscan Stud.* 73:223–28
- Töbe K, Meyer B, Fuentes V. 2010. Detection of zooplankton items in the stomach and gut content of larval krill, *Euphausia superba*, using a molecular approach. *Polar Biol.* 33:407–14
- Torres-Pratts H, Schärer MT, Schizas NV. 2009. Genetic diversity of *Chelonibia caretta*, commensal barnacles of the endangered hawksbill sea turtle *Eretmochelys imbricata* from the Caribbean (Puerto Rico). *J. Mar. Biol. Assoc. U.K.* 89:719–25
- Tsang L, Chan B, Ma K, Hsu CH, Chu K. 2007. Lack of mtDNA and morphological differentiation between two acorn barnacles *Tetraclita japonica* and *T. formosana* differing in parietes colours and geographical distribution. *Mar. Biol.* 151:147–55
- Tsang L, Chan B, Wu T, Ng W, Chatterjee T, et al. 2008. Population differentiation in the barnacle *Chthamalus malayensis*: postglacial colonization and recent connectivity across the Pacific and Indian Oceans. *Mar. Ecol. Prog. Ser.* 364:107–18
- Tsang LM, Chan BKK, Shih F-L, Chu KH, Chen CA. 2009. Host-associated speciation in the coral barnacle *Wanella milleporae* (Cirripedia: Pyrgomatidae) inhabiting the *Millepora* coral. *Mol. Ecol.* 18:1463–75
- Ueda H, Bucklin A. 2006. *Acartia (Odontacartia) obtuskai*, a new brackish-water calanoid copepod from Ariake Bay, Japan, with a redescription of the closely related *A. pacifica* from the Seto Inland Sea. *Hydrobiologia* 560:77–91
- Unal E, Frost BW, Armbrust V, Kideys AE. 2006. Phylogeography of *Calanus helgolandicus* and the Black Sea copepod *Calanus euxinus*, with notes on *Pseudocalanus elongatus* (Copepoda, Calanoida). *Deep-Sea R. II* 53:1961–75
- Undheim EAB, Norman JA, Thoen HH, Fry BG. 2010. Genetic identification of Southern Ocean octopod samples using mtCOI. *C. R. Biol.* In press

- Uthicke S, Byrne M, Conand C. 2010. Genetic barcoding of commercial Beche-de-mer species (Echinodermata: Holothuroidea). *Mol. Ecol. Resour.* 10:634–46
- Van den Broeck H, Breugelmans K, De Wolf H, Backeljau T. 2008. Completely disjunct mitochondrial DNA haplotype distribution without a phylogeographic break in a planktonic developing gastropod. *Mar. Biol.* 153:421–29
- van Helden AL, Baker AN, Dalebout ML, Reyes JC, Waerebeek KV, et al. 2002. Resurrection of *Mesoplodon traversii* (Gray, 1874), senior synonym of *M. babamondi* Reyes, Van Waerebeek, Cárdenas and Yañez, 1995 (Cetacea: Ziphiidae). *Mar. Mamm. Sci.* 18:609–21
- Vargas SM, Araujo FCF, Santos FR. 2009. DNA barcoding of Brazilian sea turtles (Testudines). *Genet. Mol. Biol.* 32:608–12
- Victor BC. 2007. *Coryphopterus kuna*, a new goby (Perciformes: Gobiidae: Gobiinae) from the western Caribbean, with the identification of the late larval stage and an estimate of the pelagic larval duration. *Zootaxa* 1526:51–61
- Victor BC. 2008. Redescription of *Coryphopterus tortugae* (Jordan) and a new allied species *Coryphopterus bol* (Perciformes: Gobiidae: Gobiinae) from the tropical western Atlantic Ocean. *J. Ocean Sci. Found.* 1:1–19
- Vilas R, Criscione C, Blouin M. 2005. A comparison between mitochondrial DNA and the ribosomal internal transcribed regions in prospecting for cryptic species of platyhelminth parasites. *Parasitology* 131:839–46
- Virgilio M, Fauvelot C, Costantini F, Abbiati M, Backeljau T. 2009. Phylogeography of the common ragworm *Hediste diversicolor* (Polychaeta: Nereididae) reveals cryptic diversity and multiple colonization events across its distribution. *Mol. Ecol.* 18:1980–94
- Vogler A, Monaghan MT. 2006. Recent advances in DNA taxonomy. *J. Zool. Syst. Evol. Res.* 45:1–10
- Vollmer S, Palumbi S. 2004. Testing the utility of internally transcribed spacer sequences in coral phylogenetics. *Mol. Ecol.* 13:2763–72
- Ward RD. 2009. DNA barcode divergence among species and genera of birds and fishes. *Mol. Ecol. Resour.* doi:10.1111/j.1755-0998.2009.02541.x
- Ward RD, Hanner R, Hebert PDN. 2009. The campaign to DNA barcode all fishes, FISH-BOL. *J. Fish Biol.* 73:1–28
- Ward RD, Holmes BH, White WT, Last PR. 2008. DNA Barcoding Australasian chondrichthyans: results and potential uses in conservation. *Mar. Freshwat. Res.* 59:57–71
- Ward RD, Holmes BH, Zemlak TS, Smith PJ. 2007. DNA barcoding discriminates spurdogs of the genus *Squalus*. In *Descriptions of New Dogfishes of the Genus Squalus (Squaloidea: Squalidae)*, ed. PR Last, WT White, JJ Pogonoski. CSIRO, Hobart, Tasmania
- Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PDN. 2005. DNA barcoding Australia's fish species. *Philos. Trans. R. Soc. Lond. B* 360:1847–57
- Wares JP, Cunningham CW. 2001. Phylogeography and historical ecology of the North Atlantic intertidal. *Evolution* 55:2455–69
- Webb KE, Barnes DKA, Clark MS, Bowden DA. 2006. DNA barcoding: A molecular tool to identify Antarctic marine larvae. *Deep-Sea Res. II* 53:1053–60
- Whitworth HC, Dawson RD, Magalon H, Baudry E. 2007. DNA barcoding cannot reliably identify species of the blowfly genus *Protocalliphora* (Diptera: Calliphoridae). *Proc. R. Soc. Lond. B* 274:1731–39
- Wiens JJ. 2007. Species delimitation: new approaches for discovering diversity. *Syst. Biol.* 56:875–78
- Will K, Rubinoff D. 2004. Myth of the molecule: DNA barcodes for species cannot replace morphology for identification and classification. *Cladistics* 20:47–55
- Williams ST, Knowlton N. 2001. Mitochondrial pseudogenes are pervasive and often insidious in the snapping shrimp genus *Alpheus*. *Mol. Biol. Evol.* 18:1484–93
- Wilson N, Hunter R, Lockhart S, Halanych K. 2007. Multiple lineages and absence of panmixia in the “circumpolar” crinoid *Promachocrinus kerguelensis* from the Atlantic sector of Antarctica. *Mar. Biol.* 152:895–904
- Wong EHK, Hanner R. 2008. DNA barcoding detects market substitution in North American seafood. *Food Res. Int.* 41:828–37
- Wörheide G. 2006. Low variation in partial cytochrome oxidase subunit I (COI) mitochondrial sequences in the coralline demosponge *Astrosclera willeyana* across the Indo-Pacific. *Mar. Biol.* 148:907–12
- Wörheide G, Erpenbeck D. 2007. DNA taxonomy of sponges—progress and perspectives. *J. Mar. Biol. Assoc. U.K.* 87:1629–33

- Xu J, Chan T-Y, Tsang LM, Chu KH. 2009. Phylogeography of the mitten crab *Eriocheir sensu stricto* in East Asia: Pleistocene isolation, population expansion and secondary contact. *Mol. Phylogen. Evol.* 52:45–56
- Yancy HF, Zemlak TS, Mason JA, Washington JD, Tenge BJ, et al. 2007. The potential use of DNA barcodes in regulatory science: applications of the Regulatory Fish Encyclopedia. *J. Food Prot.* 70:210–17
- Zemlak TS, Ward RD, Connell AD, Holmes BH, Hebert PDN. 2009. DNA barcoding reveals overlooked marine fishes. *Mol. Ecol. Resour.* 9:237–42
- Zhang AB, Sikes DS, Muster C, Li SQ. 2008. Inferring species membership using DNA sequences with back-propagation neural networks. *Syst. Biol.* 57:202–15
- Zhang J. 2010. Exploiting formalin-preserved fish specimens for resources of DNA barcoding. *Mol. Ecol. Resour.* In press; doi: 10.1111/j.1755-0998.2010.2838x
- Zimmermann J, Hajibabaei M, Blackburn DC, Hanken J, Cantin E, et al. 2008. DNA damage in preserved specimens and tissue samples: a molecular assessment. *Front. Zool.* 5:18
- Zitari-Chatti R, Chatti N, Fulgione D, Caiazza I, Aprea G, et al. 2009. Mitochondrial DNA variation in the caramote prawn *Penaeus (Melicertus) keratuburus* across a transition zone in the Mediterranean Sea. *Genetica* 136:439–47



Contents

Geologist at Sea: Aspects of Ocean History <i>Wolfgang H. Berger</i>	1
Submarine Paleoseismology Based on Turbidite Records <i>Chris Goldfinger</i>	35
Natural Processes in Delta Restoration: Application to the Mississippi Delta <i>Chris Paola, Robert R. Twilley, Douglas A. Edmonds, Wonsuck Kim, David Mohrig, Gary Parker, Enrica Viparelli, and Vaughan R. Voller</i>	67
Modeling the Dynamics of Continental Shelf Carbon <i>Eileen E. Hofmann, Bronwyn Cabill, Katja Fennel, Marjorie A.M. Friedrichs, Kimberly Hyde, Cindy Lee, Antonio Mannino, Raymond G. Najjar, John E. O'Reilly, John Wilkin, and Jianhong Xue</i>	93
Estuarine and Coastal Ocean Carbon Paradox: CO ₂ Sinks or Sites of Terrestrial Carbon Incineration? <i>Wei-Jun Cai</i>	123
Emerging Topics in Marine Methane Biogeochemistry <i>David L. Valentine</i>	147
Observations of CFCs and SF ₆ as Ocean Tracers <i>Rana A. Fine</i>	173
Nitrogen Cycle of the Open Ocean: From Genes to Ecosystems <i>Jonathan P. Zebr and Raphael M. Kudela</i>	197
Marine Primary Production in Relation to Climate Variability and Change <i>Francisco P. Chavez, Monique Messié, and J. Timothy Pennington</i>	227
Beyond the Calvin Cycle: Autotrophic Carbon Fixation in the Ocean <i>Michael Hügler and Stefan M. Sievert</i>	261
Carbon Concentrating Mechanisms in Eukaryotic Marine Phytoplankton <i>John R. Reinfelder</i>	291

Microbial Nitrogen Cycling Processes in Oxygen Minimum Zones <i>Phyllis Lam and Marcel M.M. Kuypers</i>	317
Microbial Metagenomics: Beyond the Genome <i>Jack A. Gilbert and Christopher L. Dupont</i>	347
Environmental Proteomics: Changes in the Proteome of Marine Organisms in Response to Environmental Stress, Pollutants, Infection, Symbiosis, and Development <i>Lars Tomanek</i>	373
Microbial Extracellular Enzymes and the Marine Carbon Cycle <i>Carol Arnosti</i>	401
Modeling Diverse Communities of Marine Microbes <i>Michael J. Follows and Stephanie Dutkiewicz</i>	427
Biofilms and Marine Invertebrate Larvae: What Bacteria Produce That Larvae Use to Choose Settlement Sites <i>Michael G. Hadfield</i>	453
DNA Barcoding of Marine Metazoa <i>Ann Bucklin, Dirk Steinke, and Leocadio Blanco-Bercial</i>	471
Local Adaptation in Marine Invertebrates <i>Eric Sanford and Morgan W. Kelly</i>	509
Use of Flow Cytometry to Measure Biogeochemical Rates and Processes in the Ocean <i>Michael W. Lomas, Deborah A. Bronk, and Ger van den Engh</i>	537
The Impact of Microbial Metabolism on Marine Dissolved Organic Matter <i>Elizabeth B. Kujawinski</i>	567

Errata

An online log of corrections to *Annual Review of Marine Science* articles may be found at <http://marine.annualreviews.org/errata.shtml>